

(19)



Europäisches Patentamt  
European Patent Office  
Office européen des brevets



(11)

**EP 0 882 793 A2**

(12)

**EUROPEAN PATENT APPLICATION**

(43) Date of publication:

**09.12.1998 Bulletin 1998/50**

(21) Application number: **98304208.6**

(22) Date of filing: **28.05.1998**

(51) Int Cl.<sup>6</sup>: **C12N 15/12, C12N 15/85,**

**C12N 15/11, C12N 5/10,**

**C12Q 1/02, C12Q 1/68,**

**C07K 14/705, C07K 16/28,**

**A61K 38/17, A61K 48/00,**

**G01N 33/50**

(84) Designated Contracting States:

**AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU  
MC NL PT SE**

Designated Extension States:

**AL LT LV MK RO SI**

(30) Priority: **02.06.1997 US 48240 P**

**09.12.1997 US 987289**

(71) Applicant: **SMITHKLINE BEECHAM**

**CORPORATION**

**Philadelphia Pennsylvania 19103 (US)**

(72) Inventors:

• **Hu, Erding**

**King of Prussia, Pennsylvania 19406 (US)**

• **Zhu, Yuan**

**King of Prussia, Pennsylvania 19406 (US)**

(74) Representative: **Crump, Julian Richard John et al**

**FJ Cleveland,**

**40-43 Chancery Lane**

**London WC2A 1JQ (GB)**

(54) **A human 7-tm receptor similar to murine frizzled-6 gene**

(57) hFrizzled-6 polypeptides and polynucleotides and methods for producing such polypeptides by recombinant techniques are disclosed. Also disclosed are methods for utilizing hFrizzled-6 polypeptides and polynucleotides in the design of protocols for the treatment of infections such as bacterial, fungal, protozoan and viral infections, particularly infections caused by HIV-1 or HIV-2; pain; cancers: diabetes, obesity; anorexia; bulimia; asthma; Parkinson's disease: acute heart fail-

ure; hypotension; hypertension; urinary retention; osteoporosis; angina pectoris; myocardial infarction; ulcers; asthma; allergies; benign prostatic hypertrophy; and psychotic and neurological disorders, including anxiety, schizophrenia, manic depression, delirium, dementia, severe mental retardation and dyskinesias, such as Huntington's disease or Gilles de la Tourette's syndrome, among others and diagnostic assays for such conditions.

**EP 0 882 793 A2**

**Description**

This application claims the benefit of U.S. Provisional Application No. 60/048,240, filed June 2, 1997.

**FIELD OF INVENTION**

This invention relates to newly identified polynucleotides, polypeptides encoded by them and to the use of such polynucleotides and polypeptides, and to their production. More particularly, the polynucleotides and polypeptides of the present invention relate to G-protein coupled receptor family, hereinafter referred to as hFrizzled-6. The invention also relates to inhibiting or activating the action of such polynucleotides and polypeptides.

**BACKGROUND OF THE INVENTION**

It is well established that many medically significant biological processes are mediated by proteins participating in signal transduction pathways that involve G-proteins and/or second messengers, e.g., cAMP (Lefkowitz, Nature, 1991, 351:353-354). Herein these proteins are referred to as proteins participating in pathways with G-proteins or PPG proteins. Some examples of these proteins include the GPC receptors, such as those for adrenergic agents and dopamine (Kobilka, B.K., et al., Proc. Natl Acad. Sci., USA, 1987, 84:46-50; Kobilka, B.K., et al., Science, 1987, 238:650-656; Bunzow, J.R., et al., Nature, 1988, 336:783-787), G-proteins themselves, effector proteins, e.g., phospholipase C, adenylyl cyclase, and phosphodiesterase, and actuator proteins, e.g., protein kinase A and protein kinase C (Simon, M. I., et al., Science, 1991, 252:802-8).

For example, in one form of signal transduction, the effect of hormone binding is activation of the enzyme, adenylyl cyclase, inside the cell. Enzyme activation by hormones is dependent on the presence of the nucleotide, GTP. GTP also influences hormone binding. A G-protein connects the hormone receptor to adenylyl cyclase. G-protein was shown to exchange GTP for bound GDP when activated by a hormone receptor. The GTP-carrying form then binds to activated adenylyl cyclase. Hydrolysis of GTP to GDP, catalyzed by the G-protein itself, returns the G-protein to its basal, inactive form. Thus, the G-protein serves a dual role, as an intermediate that relays the signal from receptor to effector, and as a clock that controls the duration of the signal.

The membrane protein gene superfamily of G-protein coupled receptors has been characterized as having seven putative transmembrane domains. The domains are believed to represent transmembrane  $\alpha$ -helices connected by extracellular or cytoplasmic loops. G-protein coupled receptors include a wide range of biologically active receptors, such as hormone, viral, growth factor and neuroreceptors.

G-protein coupled receptors (otherwise known as 7TM receptors) have been characterized as including these seven conserved hydrophobic stretches of about 20 to 30 amino acids, connecting at least eight divergent hydrophilic loops. The G-protein family of coupled receptors includes dopamine receptors which bind to neuroleptic drugs used for treating psychotic and neurological disorders. Other examples of members of this family include, but are not limited to, calcitonin, adrenergic, endothelin, cAMP, adenosine, muscarinic, acetylcholine, serotonin, histamine, thrombin, kinin, follicle stimulating hormone, opsins, endothelial differentiation gene-1, rhodopsins, odorant, and cytomegalovirus receptors.

Most G-protein coupled receptors have single conserved cysteine residues in each of the first two extracellular loops which form disulfide bonds that are believed to stabilize functional protein structure. The 7 transmembrane regions are designated as TM1, TM2, TM3, TM4, TM5, TM6, and TM7. TM3 has been implicated in signal transduction.

Phosphorylation and lipidation (palmitoylation or farnesylation) of cysteine residues can influence signal transduction of some G-protein coupled receptors. Most G-protein coupled receptors contain potential phosphorylation sites within the third cytoplasmic loop and/or the carboxy terminus. For several G-protein coupled receptors, such as the  $\beta$ -adrenoreceptor, phosphorylation by protein kinase A and/or specific receptor kinases mediates receptor desensitization.

For some receptors, the ligand binding sites of G-protein coupled receptors are believed to comprise hydrophilic sockets formed by several G-protein coupled receptor transmembrane domains, said sockets being surrounded by hydrophobic residues of the G-protein coupled receptors. The hydrophilic side of each G-protein coupled receptor transmembrane helix is postulated to face inward and form a polar ligand binding site. TM3 has been implicated in several G-protein coupled receptors as having a ligand binding site, such as the TM3 aspartate residue. TM5 serines, a TM6 asparagine and TM6 or TM7 phenylalanines or tyrosines are also implicated in ligand binding.

G-protein coupled receptors can be intracellularly coupled by heterotrimeric G-proteins to various intracellular enzymes, ion channels and transporters (see, Johnson et al., Endoc. Rev., 1989, 10:317-331). Different G-protein  $\alpha$ -subunits preferentially stimulate particular effectors to modulate various biological functions in a cell. Phosphorylation of cytoplasmic residues of G-protein coupled receptors has been identified as an important mechanism for the regulation of G-protein coupling of some G-protein coupled receptors. G-protein coupled receptors are found in numerous

sites within a mammalian host. Over the past 15 years, nearly 350 therapeutic agents targeting 7 transmembrane (7 TM) receptors have been successfully introduced onto the market.

This indicates that these receptors have an established, proven history as therapeutic targets. Clearly there is a need for identification and characterization of further receptors which can play a role in preventing, ameliorating or correcting dysfunctions or diseases, including, but not limited to, infections such as bacterial, fungal, protozoan and viral infections, particularly infections caused by HIV-1 or HIV-2; pain; cancers; diabetes; obesity; anorexia; bulimia; asthma; Parkinson's disease; acute heart failure; hypotension; hypertension; urinary retention; osteoporosis; angina pectoris; myocardial infarction; ulcers; asthma; allergies; benign prostatic hypertrophy; and psychotic and neurological disorders, including anxiety, schizophrenia, manic depression, delirium, dementia, severe mental retardation and dyskinesias, such as Huntington's disease or Gilles de la Tourette's syndrome.

## SUMMARY OF THE INVENTION

In one aspect, the invention relates to hFrizzled-6 polypeptides and recombinant materials and methods for their production. Another aspect of the invention relates to methods for using such hFrizzled-6 polypeptides and polynucleotides. Such uses include the treatment of infections such as bacterial, fungal, protozoan and viral infections, particularly infections caused by HIV-1 or HIV-2; pain; cancers; diabetes; obesity; anorexia; bulimia; asthma; Parkinson's disease; acute heart failure; hypotension; hypertension; urinary retention; osteoporosis; angina pectoris; myocardial infarction; ulcers; asthma; allergies; benign prostatic hypertrophy; and psychotic and neurological disorders, including anxiety, schizophrenia, manic depression, delirium, dementia, severe mental retardation and dyskinesias, such as Huntington's disease or Gilles de la Tourette's syndrome, among others. In still another aspect, the invention relates to methods to identify agonists and antagonists using the materials provided by the invention, and treating conditions associated with hFrizzled-6 imbalance with the identified compounds. Yet another aspect of the invention relates to diagnostic assays for detecting diseases associated with inappropriate hFrizzled-6 activity or levels.

## DESCRIPTION OF THE INVENTION

### Definitions

The following definitions are provided to facilitate understanding of certain terms used frequently herein.

"hFrizzled-6" refers, among others, to a polypeptide comprising the amino acid sequence set forth in SEQ ID NO: 2, or an allelic variant thereof.

"Receptor Activity" or "Biological Activity of the Receptor" refers to the metabolic or physiologic function of said hFrizzled-6 including similar activities or improved activities or these activities with decreased undesirable side-effects. Also included are antigenic and immunogenic activities of said hFrizzled-6.

"hFrizzled-6 gene" refers to a polynucleotide comprising the nucleotide sequence set forth in SEQ ID NO: 1 or allelic variants thereof and/or their complements.

"Antibodies" as used herein includes polyclonal and monoclonal antibodies, chimeric, single chain, and humanized antibodies, as well as Fab fragments, including the products of an Fab or other immunoglobulin expression library.

"Isolated" means altered "by the hand of man" from the natural state. If an "isolated" composition or substance occurs in nature, it has been changed or removed from its original environment, or both. For example, a polynucleotide or a polypeptide naturally present in a living animal is not "isolated," but the same polynucleotide or polypeptide separated from the coexisting materials of its natural state is "isolated", as the term is employed herein.

"Polynucleotide" generally refers to any polyribonucleotide or polydeoxiribonucleotide, which may be unmodified RNA or DNA or modified RNA or DNA. "Polynucleotides" include, without limitation single- and double-stranded DNA, DNA that is a mixture of single- and double-stranded regions, single- and double-stranded RNA, and RNA that is mixture of single- and double-stranded regions, hybrid molecules comprising DNA and RNA that may be single-stranded or, more typically, double-stranded or a mixture of single- and double-stranded regions. In addition, "polynucleotide" refers to triple-stranded regions comprising RNA or DNA or both RNA and DNA. The term polynucleotide also includes DNAs or RNAs containing one or more modified bases and DNAs or RNAs with backbones modified for stability or for other reasons. "Modified" bases include, for example, tritylated bases and unusual bases such as inosine. A variety of modifications has been made to DNA and RNA; thus, "polynucleotide" embraces chemically, enzymatically or metabolically modified forms of polynucleotides as typically found in nature, as well as the chemical forms of DNA and RNA characteristic of viruses and cells. "Polynucleotide" also embraces relatively short polynucleotides, often referred to as oligonucleotides.

"Polypeptide" refers to any peptide or protein comprising two or more amino acids joined to each other by peptide bonds or modified peptide bonds, i.e., peptide isosteres. "Polypeptide" refers to both short chains, commonly referred to as peptides, oligopeptides or oligomers, and to longer chains, generally referred to as proteins. Polypeptides may

contain amino acids other than the 20 gene-encoded amino acids. "Polypeptides" include amino acid sequences modified either by natural processes, such as posttranslational processing, or by chemical modification techniques which are well known in the art. Such modifications are well described in basic texts and in more detailed monographs, as well as in a voluminous research literature. Modifications can occur anywhere in a polypeptide, including the peptide backbone, the amino acid side-chains and the amino or carboxyl termini. It will be appreciated that the same type of modification may be present in the same or varying degrees at several sites in a given polypeptide. Also, a given polypeptide may contain many types of modifications. Polypeptides may be branched as a result of ubiquitination, and they may be cyclic, with or without branching. Cyclic, branched and branched cyclic polypeptides may result from posttranslation natural processes or may be made by synthetic methods. Modifications include acetylation, acylation, ADP-ribosylation, amidation, covalent attachment of flavin, covalent attachment of a heme moiety, covalent attachment of a nucleotide or nucleotide derivative, covalent attachment of a lipid or lipid derivative, covalent attachment of phosphatidylinositol, cross-linking, cyclization, disulfide bond formation, demethylation, formation of covalent cross-links, formation of cystine, formation of pyroglutamate, formylation, gamma-carboxylation, glycosylation, GPI anchor formation, hydroxylation, iodination, methylation, myristoylation, oxidation, proteolytic processing, phosphorylation, prenylation, racemization, selenoylation, sulfation, transfer-RNA mediated addition of amino acids to proteins such as arginylation, and ubiquitination. See, for instance, PROTEINS - STRUCTURE AND MOLECULAR PROPERTIES, 2nd Ed., T. E. Creighton, W. H. Freeman and Company, New York, 1993 and Wold, F., Posttranslational Protein Modifications: Perspectives and Prospects, pgs. 1-12 in POSTTRANSLATIONAL COVALENT MODIFICATION OF PROTEINS, B. C. Johnson, Ed., Academic Press, New York, 1983; Seifter *et al.*, "Analysis for protein modifications and nonprotein cofactors", *Meth Enzymol* (1990) 182:626-646 and Rattan *et al.*, "Protein Synthesis: Posttranslational Modifications and Aging", *Ann NY Acad Sci* (1992) 663:48-62.

"Variant" as the term is used herein, is a polynucleotide or polypeptide that differs from a reference polynucleotide or polypeptide respectively, but retains essential properties. A typical variant of a polynucleotide differs in nucleotide sequence from another, reference polynucleotide. Changes in the nucleotide sequence of the variant may or may not alter the amino acid sequence of a polypeptide encoded by the reference polynucleotide. Nucleotide changes may result in amino acid substitutions, additions, deletions, fusions and truncations in the polypeptide encoded by the reference sequence, as discussed below. A typical variant of a polypeptide differs in amino acid sequence from another, reference polypeptide. Generally, differences are limited so that the sequences of the reference polypeptide and the variant are closely similar overall and, in many regions, identical. A variant and reference polypeptide may differ in amino acid sequence by one or more substitutions, additions, deletions in any combination. A substituted or inserted amino acid residue may or may not be one encoded by the genetic code. A variant of a polynucleotide or polypeptide may be a naturally occurring such as an allelic variant, or it may be a variant that is not known to occur naturally. Non-naturally occurring variants of polynucleotides and polypeptides may be made by mutagenesis techniques or by direct synthesis.

"Identity" is a measure of the identity of nucleotide sequences or amino acid sequences. In general, the sequences are aligned so that the highest order match is obtained. "Identity" *per se* has an art-recognized meaning and can be calculated using published techniques. See, e.g.: (COMPUTATIONAL MOLECULAR BIOLOGY, Lesk, A.M., ed., Oxford University Press, New York, 1988; BIOCOMPUTING: INFORMATICS AND GENOME PROJECTS, Smith, D.W., ed., Academic Press, New York, 1993; COMPUTER ANALYSIS OF SEQUENCE DATA, PART I, Griffin, A.M., and Griffin, H.G., eds., Humana Press, New Jersey, 1994; SEQUENCE ANALYSIS IN MOLECULAR BIOLOGY, von Heinje, G., Academic Press, 1987; and SEQUENCE ANALYSIS PRIMER, Gribskov, M. and Devereux, J., eds., M Stockton Press, New York, 1991). While there exist a number of methods to measure identity between two polynucleotide or polypeptide sequences, the term "identity" is well known to skilled artisans (Carillo, H., and Lipton, D., *SIAM J Applied Math* (1988) 48:1073). Methods commonly employed to determine identity or similarity between two sequences include, but are not limited to, those disclosed in Guide to Huge Computers, Martin J. Bishop, ed., Academic Press, San Diego, 1994, and Carillo, H., and Lipton, D., *SIAM J Applied Math* (1988) 48:1073. Methods to determine identity and similarity are codified in computer programs. Preferred computer program methods to determine identity and similarity between two sequences include, but are not limited to, GCS program package (Devereux, J., *et al.*, *Nucleic Acids Research* (1984) 12(1):387), BLASTP, BLASTN, FASTA (Atschul, S.F. *et al.*, *J Molec Biol* (1990) 215:403).

As an illustration, by a polynucleotide having a nucleotide sequence having at least, for example, 95% "identity" to a reference nucleotide sequence of SEQ ID NO: 1 is intended that the nucleotide sequence of the polynucleotide is identical to the reference sequence except that the polynucleotide sequence may include up to five point mutations per each 100 nucleotides of the reference nucleotide sequence of SEQ ID NO: 1. In other words, to obtain a polynucleotide having a nucleotide sequence at least 95% identical to a reference nucleotide sequence, up to 5% of the nucleotides in the reference sequence may be deleted or substituted with another nucleotide, or a number of nucleotides up to 5% of the total nucleotides in the reference sequence may be inserted into the reference sequence. These mutations of the reference sequence may occur at the 5' or 3' terminal positions of the reference nucleotide sequence or anywhere between those terminal positions, interspersed either individually among nucleotides in the reference

sequence or in one or more contiguous groups within the reference sequence.

Similarly, by a polypeptide having an amino acid sequence having at least, for example, 95% "identity" to a reference amino acid sequence of SEQ ID NO:2 is intended that the amino acid sequence of the polypeptide is identical to the reference sequence except that the polypeptide sequence may include up to five amino acid alterations per each 100 amino acids of the reference amino acid of SEQ ID NO: 2. In other words, to obtain a polypeptide having an amino acid sequence at least 95% identical to a reference amino acid sequence, up to 5% of the amino acid residues in the reference sequence may be deleted or substituted with another amino acid, or a number of amino acids up to 5% of the total amino acid residues in the reference sequence may be inserted into the reference sequence. These alterations of the reference sequence may occur at the amino or carboxy terminal positions of the reference amino acid sequence or anywhere between those terminal positions, interspersed either individually among residues in the reference sequence or in one or more contiguous groups within the reference sequence.

### Polypeptides of the Invention

In one aspect, the present invention relates to hFrizzled-6 polypeptides (or hFrizzled-6 proteins). The hFrizzled-6 polypeptides include the polypeptides of SEQ ID NOS:2 and 4; as well as polypeptides comprising the amino acid sequence of SEQ ID NO:2; and polypeptides comprising the amino acid sequence which have at least 80% identity to that of SEQ ID NO:2 over its entire length, and still more preferably at least 90% identity, and even still more preferably at least 95% identity to SEQ ID NO: 2. Furthermore, those with at least 97-99% are highly preferred. Also included within hFrizzled-6 polypeptides are polypeptides having the amino acid sequence which have at least 80% identity to the polypeptide having the amino acid sequence of SEQ ID NO: 2 over its entire length, and still more preferably at least 90% identity, and even still more preferably at least 95% identity to SEQ ID NO: 2. Furthermore, those with at least 97-99% are highly preferred. Preferably hFrizzled-6 polypeptides exhibit at least one biological activity of the receptor.

The hFrizzled-6 polypeptides may be in the form of the "mature" protein or may be a part of a larger protein such as a fusion protein. It is often advantageous to include an additional amino acid sequence which contains secretory or leader sequences, pro-sequences, sequences which aid in purification such as multiple histidine residues, or an additional sequence for stability during recombinant production.

Fragments of the hFrizzled-6 polypeptides are also included in the invention. A fragment is a polypeptide having an amino acid sequence that entirely is the same as part, but not all, of the amino acid sequence of the aforementioned hFrizzled-6 polypeptides. As with hFrizzled-6 polypeptides, fragments may be "free-standing," or comprised within a larger polypeptide of which they form a part or region, most preferably as a single continuous region. Representative examples of polypeptide fragments of the invention, include, for example, fragments from about amino acid number 1-20, 21-40, 41-60, 61-80, 81-100, and 101 to the end of hFrizzled-6 polypeptide. In this context "about" includes the particularly recited ranges larger or smaller by several, 5, 4, 3, 2 or 1 amino acid at either extreme or at both extremes.

Preferred fragments include, for example, truncation polypeptides having the amino acid sequence of hFrizzled-6 polypeptides, except for deletion of a continuous series of residues that includes the amino terminus, or a continuous series of residues that includes the carboxyl terminus or deletion of two continuous series of residues, one including the amino terminus and one including the carboxyl terminus. Also preferred are fragments characterized by structural or functional attributes such as fragments that comprise alpha-helix and alpha-helix forming regions, beta-sheet and beta-sheet-forming regions, turn and turn-forming regions, coil and coil-forming regions, hydrophilic regions, hydrophobic regions, alpha amphipathic regions, beta amphipathic regions, flexible regions, surface-forming regions, substrate binding region, and high antigenic index regions. Other preferred fragments are biologically active fragments. Biologically active fragments are those that mediate receptor activity, including those with a similar activity or an improved activity, or with a decreased undesirable activity. Also included are those that are antigenic or immunogenic in an animal, especially in a human.

Preferably, all of these polypeptide fragments retain the biological activity of the receptor, including antigenic activity. Among the most preferred fragment is that having the amino acid sequence of SEQ ID NO: 4. Variants of the defined sequence and fragments also form part of the present invention. Preferred variants are those that vary from the referents by conservative amino acid substitutions -- i.e., those that substitute a residue with another of like characteristics. Typical such substitutions are among Ala, Val, Leu and Ile; among Ser and Thr; among the acidic residues Asp and Glu; among Asn and Gln; and among the basic residues Lys and Arg; or aromatic residues Phe and Tyr. Particularly preferred are variants in which several, 5-10, 1-5, or 1-2 amino acids are substituted, deleted, or added in any combination.

The hFrizzled-6 polypeptides of the invention can be prepared in any suitable manner. Such polypeptides include isolated naturally occurring polypeptides, recombinantly produced polypeptides, synthetically produced polypeptides, or polypeptides produced by a combination of these methods. Means for preparing such polypeptides are well understood in the art.

**Polynucleotides of the Invention**

Another aspect of the invention relates to hFrizzled-6 polynucleotides. hFrizzled-6 polynucleotides include isolated polynucleotides which encode the hFrizzled-6 polypeptides and fragments, and polynucleotides closely related thereto. More specifically, hFrizzled-6 polynucleotide of the invention include a polynucleotide comprising the nucleotide sequence contained in SEQ ID NO:1 encoding a hFrizzled-6 polypeptide of SEQ ID NO: 2, and polynucleotides having the particular sequences of SEQ ID NOS:1 and 3. hFrizzled-6 polynucleotides further include a polynucleotide comprising a nucleotide sequence that has at least 80% identity over its entire length to a nucleotide sequence encoding the hFrizzled-6 polypeptide of SEQ ID NO:2, and a polynucleotide comprising a nucleotide sequence that is at least 80% identical to that of SEQ ID NO: 1 over its entire length. In this regard, polynucleotides at least 90% identical are particularly preferred, and those with at least 95% are especially preferred. Furthermore, those with at least 97% are highly preferred and those with at least 98-99% are most highly preferred, with at least 99% being the most preferred. Also included under hFrizzled-6 polynucleotides are a nucleotide sequence which has sufficient identity to a nucleotide sequence contained in SEQ ID NO:1 to hybridize under conditions useable for amplification or for use as a probe or marker. The invention also provides polynucleotides which are complementary to such hFrizzled-6 polynucleotides.

hFrizzled-6 of the invention is structurally related to other proteins of the G-protein coupled receptor family, as shown by the results of sequencing the cDNA encoding human hFrizzled-6. The cDNA sequence of SEQ ID NO:1 contains an open reading frame (nucleotide number 134 to 2252) encoding a polypeptide of 706 amino acids of SEQ ID NO:2. The amino acid sequence of Table 1 (SEQ ID NO:2) has about 67.3% identity (using fast-A) in 706 amino acid residues with mouse transmembrane receptor frizzled-6 [Wang, Y., Macke, J. P., Abella, B. S., Andreasson, K., Worley, P., Gilbert, D. J., Copeland, N. G., Jenkins, N. A. and Nathans, J.; J. Biol. Chem. 271(8), 4468-4476 (1996)]. Furthermore, the amino acid sequence of Table 1 (SEQ ID NO:2) shares about 52% identity over 706 amino acid residues with mouse Frizzled-3 [Wang, Y., Macke, J. P., Abella, B. S., Andreasson, K., Worley, P., Gilbert, D. J., Copeland, N. G., Jenkins, N. A. and Nathans, J.; J. Biol. Chem. 271 (8), 4468-4476 (1996)]. The nucleotide sequence of Table 1 (SEQ ID NO:1) has about 65% identity (using Fast-N) in 2880 nucleotide residues with mouse Frizzled-6 receptor [Wang, Y., Macke, J. P., Abella, B. S., Andreasson, K., Worley, P., Gilbert, D. J., Copeland, N. G., Jenkins, N. A. and Nathans, J.; J. Biol. Chem. 271 (8), 4468-4476 (1996)]. Furthermore, the nucleotide sequence of Table 1 (SEQ ID NO:1) has about 63% identity over 2880 nucleotide residues with mouse frizzled-3 [Wang, Y., Macke, J. P., Abella, B. S., Andreasson, K., Worley, P., Gilbert, D. J., Copeland, N. G., Jenkins, N. A. and Nathans, J.; J. Biol. Chem. 271 (8), 4468-4476 (1996)]. Thus, hFrizzled-6 polypeptides and polynucleotides of the present invention are expected to have, inter alia, similar biological functions/properties to their homologous polypeptides and polynucleotides, and their utility is obvious to anyone skilled in the art.

**Table 1<sup>a</sup>**

CGGAGCCGTCTCAGGTCCCTGGGGGAACGGTGGGTTAGACGGGGACGGGAAGGGACAGCGGCCTTCGA  
 CCGCCCCCGAGTAATTGACCCAGGACTCATTTTCAGGAAAGCCTGAAAATGAGTAAAATAGTGAAATG  
 AGGAATTTGAACATTTTATCTTTGGATGGGGATCTTCTGAGGATGCAAAGAGTGATTTCATCCAAGCCAT  
 GTGGTAAAATCAGGAATTTGAAGAAAATGGAGATGTTTACATTTTTTGTGACGTGTATTTTTCTACCCC  
 TCCTAAGAGGGCACAGTCTCTTCACCTGTGAACCAATTACTGTTCCAGATGTATGAAAATGGCCTACA  
 ACATGACGTTTTTCCCTAATCTGATGGGTCAATTATGACCAGAgTATTGCCGCGGTGGAAATGGAgCATT  
 TTCTTCCTCTCGCAAATCTGGAATGTTACCAAACATTGAAACTTTCTCTGCAAAGCATTGTGTACCAA  
 CCTGCATAgAACAAATTCATGTGGTTCACCTTGTCgTAAACTTTGTGAGAAAGTATATTcTGAttgCa  
 aAAAATTAATTGACACTTTTGGGATCCGATGGCCTGAGGAGCTTGAATGTGACAGATTACAATACTGTG  
 ATGAGACTGTTTCTGTAACTTTTGATCCACACACAGAATTTCTTGGTCCTCAGAAGGAAACAGAACAG  
 TCCAAAGAGACATTGGATTTTGGTGTCCAAGGCATCTTAAGACTTCTGGGGGACAAGGATATAAgTtC  
 TgGGAATTGACCAgTGTGCGCCTCCATGcCCCAACATGTaTTTTAAAAGTGATGAGCTAgAGTTTGCAA  
 AAAGTTTtATTGGAACAGTTTCAATATTTTGTCTTTGTGCAACTCTGTTTACATTCCTTACTTTTTTAA  
 TTGATGTTAGAAGATTAGATACCCAgAGAGACCAATTATATATTACTCTGTCTGTTACAGCATTGTAT  
 CTCTTATGTACTTCATTGGATTTTGGTGGCGATAGCACAGCCTGCAATAAGGCAGATGAGAAGCTAG  
 AACTTGGTGACACTGTTGTCTAGGCTCTCAAAATAAGGCTTGCACCGTTTTGTTTCATGCTTTTGTATT  
 TTTTCACAATGGCTGGCACTGTGTGGTGGGTGATTCTTACCATTACTTGGTTCTTAgCTGCAGGAAGAA  
 AATGGAGTTGTGAAGCCATCGAGCAAAAAGCAGTGTGGTTTCATGCTGTTGCATGGGGAACACCAaGTT  
 TCCTGACTGTTATGCTTCTTGCTCTGAACAAAGTTGAAGGAGACAACATTAGTGGAGTTcGCTTTGTTG  
 GCCTTTATGACCTGGATGCTTCTCGCTACTTTGTACTCTTGCCACTGTGCCTTTGTGTGTTTGTGGGC  
 TCTCTCTTCTTTTAGCTGGCATTATTTCTTAAATCATGTTTCGACAAGTCATACAACATGATGGCCGGA  
 ACCAAGAAAACTAAAGAAATTTATGATTTCGAATTGGAGTCTTCAGCGGCTTGTATCTTGTGCCATTAG  
 TGACACTTCTCGGATGTTACGTCTATGAgCAAGTGAACAGGATTACCTGGGAGATAACTTGGGTCTCTG  
 ATCATTGTCGTCAGTACCATATCCCATGTCCTTATCAGGCAAAAGCAAAAGCTCGACCAGAATTGGCTT  
 TATTTATGATAAAATACCTGATGACATTAATTGTTGGCATCTCTGCTGTCTTCTGGGTGGAAgCAAAA  
 AGACATGCACAGAATGGGctGGGTTTTTTAAacGAAATCGCGGGAGAGATCCAATCAGTGAAAGTCGAA  
 GAGTACTACAGGAATCATGTGAGTtTTTCTTAAAGCACAAATTCTAAAGTTAAACACAAAAGAAGCACT  
 ATAAACCGAGTTCACgCAAGCTGAAGGTCAATTCCaAATCCATGGGAACCAgCACAGGAGCTACaGCAA

aTCATGGCACTTCTGCAGTAGCAATTACTAgCCaTGATTACCTAGGACAAGAACTTTGACAGAAATCC  
 AAACCTCACCAGAAACATCAATGAGAGAGGTGAAAGCGGACGGAGCTAGCACCCCCAGGTTAAGAGAAC  
 5 AGGACTGTGGTGAACCTGCCTCGCCAGCAGCATCCATCTCCAGACTCTCTGGGGAACAGGTCGACGGGA  
 AGGGCCAGGCAGGCAGTGTATCTGAAAGTGC GCGGAgTGAAGGAAGGATTAgTCCaaAGAGTGATATTA  
 CTGACACTGGCCTGGCACAGAgCAACAATTTGCAGGTCCCCAGTTCTTCAgAACCAAGCAGCCTCaAAG  
 10 GTTCCACATCTCTGCTTGTTCACCCAGTTTCAGGAGTGAGAAAAGAGCAGGGAGGTGGTTGTCATTACAG  
 ATACTTGAAGAACATTTTCTCTCGTTACTCAGAAGCAAATTTGTGTTACACTGGAAGTGACCTATGCAC  
 TGTTTTGTAAGAATCACTGTTACGTTCTTCTTTTGCACTTAAAGTTGCATTGCCTACTGTTATACTGGA  
 AAAAAATAGAGTTCAAGAATAATATGACTCATTTACACAAAGGTTAATGACAACAATATACCTGAAAAC  
 15 AGAAATGTGCAGGTTAATAATATTTTTTTAATAGTGTGGGAGGACAGAGTTAGAGGAATCTTCCTTTTC  
 TATTTATGAAGATTCTACTCTTGGTAAGAGTATTTTAAGATGTACTATGCTATTTTACTTTTTTGATAT  
 AAAATCAAGATATTTCTTTGCTGAAGTATTTAAATCTTATCCTTGATCTTTTTATACATATTTGAAAA  
 20 TAAGCTTATATGTATTTGAACTTTTTTGAAATCCTATTCCAGTaTTTTTATCATGCTATTGTGATATtT  
 TAGCACTTTGGTAGCTTTtACACTGAAATTTCTAAGAAAATTGTAAATAG

<sup>a</sup> A nucleotide sequence of a human hFrizzled-6 (SEQ ID NO: 1).

**Table 2<sup>b</sup>**

MEMFTFLLTClFLPLLRGHSLFTCEPITVPRCMKMayNMTFFPNLMGHYDQSIAAVEMEHFLPLANLEC  
 30 SPNIETFLCKAFVPTCIEQIHVVPPCRKLCEKVYS DCKKLIDTFGIRWPEELEC DRLQYCD ETVPVTFD  
 PHTEFLGPGKETE QVQRDIGFWCPRHLKTS GGQGYKFLGIDQCAPP CPNMYFKSDELEFAKSF IGTVSI  
 FCLCATLFTFLTFLIDVRRFRYPERPIIYYSVCYSIVSLMYFIGFLLGDSTACNKAD EKL ELGDTVVLG  
 35 SQNKACTVLFMLLYFFT MAGTVWV VIL TITWFLAAGR KWSCEAIEQKAVWFHAVA WGTPSFLTVM LLLAL  
 NKVEGDNISGVRFVGLYDL DASRYFVLLPLCLCFVGLS LLLAGIISLNHVRQVIQH DGRNQEK LKKFM  
 IRIGVFSGLYLVPLVTL LGCYVYEQVNRITWEITWSDHCRQYHIPC PYQAKAKARPELALFMI KYLMT  
 LIVGISAVFWVGS KKTCTEWAGFFKRN RGRDPISESRRVLQESCEFFLKHNSKV KHKKKHYK PSSRKLK  
 40 VISKSMGTSTGATANHGTS AVAITSHDYLGQETL TEIQTS PETSMREVKADGASTPRLREQDCGE PASP  
 AASISRLSGEQVDGKGQAGSVSESARSEGRISP KSDITDTGLAQSNLQVPSSSEPSSLKGSTSL LVHP  
 VSGVRKEQGGGCHSDT

<sup>b</sup> An amino acid sequence of a human hFrizzled-6 (SEQ ID NO: 2).

One polynucleotide of the present invention encoding hFrizzled-6 may be obtained using standard cloning and screening, from a cDNA library derived from mRNA in cells of human thyroid using the expressed sequence tag (EST) analysis (Adams, M.D., *et al. Science* (1991) 252:1651-1656; Adams, M.D. *et al., Nature*, (1992) 355:632-634; Adams, M.D., *et al., Nature* (1995) 377 Supp:3-174). Polynucleotides of the invention can also be obtained from natural sources such as genomic DNA libraries or can be synthesized using well known and commercially available techniques.

The nucleotide sequence encoding hFrizzled-6 polypeptide of SEQ ID NO:2 may be identical to the polypeptide encoding sequence contained in Table 1 (nucleotide number 134 to 2252 of SEQ ID NO: 1), or it may be a sequence, which as a result of the redundancy (degeneracy) of the genetic code, also encodes the polypeptide of SEQ ID NO:2.

When the polynucleotides of the invention are used for the recombinant production of hFrizzled-6 polypeptide, the polynucleotide may include the coding sequence for the mature polypeptide or a fragment thereof, by itself; the coding sequence for the mature polypeptide or fragment in reading frame with other coding sequences, such as those encoding a leader or secretory sequence, a pre-, or pro- or prepro- protein sequence, or other fusion peptide portions. For



example, a marker sequence which facilitates purification of the fused polypeptide can be encoded. In certain preferred embodiments of this aspect of the invention, the marker sequence is a hexa-histidine peptide, as provided in the pQE vector (Qiagen, Inc.) and described in Gentz *et al.*, *Proc Natl Acad Sci USA* (1989) 86:821-824, or is an HA tag. The polynucleotide may also contain non-coding 5' and 3' sequences, such as transcribed, non-translated sequences, splicing and polyadenylation signals, ribosome binding sites and sequences that stabilize mRNA.

Further preferred embodiments are polynucleotides encoding hFrizzled-6 variants comprising the amino acid sequence of hFrizzled-6 polypeptide of Table 2 (SEQ ID NO:2) in which several, 5-10, 1-5, 1-3, 1-2 or 1 amino acid residues are substituted, deleted or added, in any combination. Among the preferred polynucleotides of the present invention is contained in Table 3 (SEQ ID NO: 3) encoding the amino acid sequence of Table 4 (SEQ ID NO: 4).

**Table 3<sup>c</sup>**

```
TCTGAACAAAGTTGAAGGAGACAACATTAGTGGAGTTGCTTTGTTGGCCTTTATGACCT
GGATGCTTCTCGCTACTTTGTACTCTTGCCACTGTGCCTTTNTGTGTTTGTGGGCTCTC
TCTTCTTTTAGCTGGCATTATTTCCCTAAATCATGTTGACAAAGTCATACAACATGATGG
CCGGAACCAAGAAAACTAAAGAAATTTATGATTCGAATTGGAGTCTTCAGCGGCTTTGT
ATCTTGTGCCATTAGTGACACTTCTCGGATGTTACGTCTATGAGCAAGTGAACAGGATTA
CCTGGGAGATACTT
```

<sup>c</sup> A partial nucleotide sequence of a human hFrizzled-6 (SEQ ID NO: 3).

**Table 4<sup>d</sup>**

```
LNKVEGDNISGVCFVGLYDLDSRYF
```

<sup>d</sup> A partial amino acid sequence of a human hFrizzled-6 (SEQ ID NO: 4).

The present invention further relates to polynucleotides that hybridize to the herein above-described sequences. In this regard, the present invention especially relates to polynucleotides which hybridize under stringent conditions to the herein above-described polynucleotides. As herein used, the term "stringent conditions" means hybridization will occur only if there is at least 80%, and preferably at least 90%, and more preferably at least 95%, yet even more preferably 97-99% identity between the sequences.

Polynucleotides of the invention, which are identical or sufficiently identical to a nucleotide sequence contained in SEQ ID NO: 1 or a fragment thereof, may be used as hybridization probes for cDNA and genomic DNA, to isolate full-length cDNAs and genomic clones encoding hFrizzled-6 and to isolate cDNA and genomic clones of other genes (including genes encoding homologs and orthologs from species other than human) that have a high sequence similarity to the hFrizzled-6 gene. Such hybridization techniques are known to those of skill in the art. Typically these nucleotide sequences are 80% identical, preferably 90% identical, more preferably 95% identical to that of the referent. The probes generally will comprise at least 15 nucleotides. Preferably, such probes will have at least 30 nucleotides and may have at least 50 nucleotides. Particularly preferred probes will range between 30 and 50 nucleotides.

In one embodiment, to obtain a polynucleotide encoding hFrizzled-6 polypeptide, including homologs and orthologs from species other than human, comprises the steps of screening an appropriate library under stringent hybridization conditions with a labeled probe having the SEQ ID NO: 1 or a fragment thereof (including that of SEQ ID NO: 3), and isolating full-length cDNA and genomic clones containing said polynucleotide sequence. Such hybridization techniques are well known to those of skill in the art. Stringent hybridization conditions are as defined above or alternatively conditions under overnight incubation at 42°C in a solution comprising: 50% formamide, 5xSSC (150mM NaCl, 15mM trisodium citrate), 50 mM sodium phosphate (pH7.6), 5x Denhardt's solution, 10 % dextran sulfate, and 20 microgram/ml denatured, sheared salmon sperm DNA, followed by washing the filters in 0.1x SSC at about 65°C.

The polynucleotides and polypeptides of the present invention may be employed as research reagents and materials for discovery of treatments and diagnostics to animal and human disease.

## Vectors, Host Cells, Expression

The present invention also relates to vectors which comprise a polynucleotide or polynucleotides of the present invention, and host cells which are genetically engineered with vectors of the invention and to the production of polypep-

tides of the invention by recombinant techniques. Cell-free translation systems can also be employed to produce such proteins using RNAs derived from the DNA constructs of the present invention.

For recombinant production, host cells can be genetically engineered to incorporate expression systems or portions thereof for polynucleotides of the present invention. Introduction of polynucleotides into host cells can be effected by methods described in many standard laboratory manuals, such as Davis et al., *BASIC METHODS IN MOLECULAR BIOLOGY* (1986) and Sambrook et al., *MOLECULAR CLONING: A LABORATORY MANUAL*, 2nd Ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1989) such as calcium phosphate transfection, DEAF-dextran mediated transfection, transvection, microinjection, cationic lipid-mediated transfection, electroporation, transduction, scrape loading, ballistic introduction or infection.

Representative examples of appropriate hosts include bacterial cells, such as streptococci, staphylococci, *E. coli*, *Streptomyces* and *Bacillus subtilis* cells; fungal cells, such as yeast cells and *Aspergillus* cells; insect cells such as *Drosophila* S2 and *Spodoptera* Sf9 cells; animal cells such as CHO, COS, HeLa, C127, 3T3, BHK, HEK 293 and Bowes melanoma cells; and plant cells.

A great variety of expression systems can be used. Such systems include, among others, chromosomal, episomal and virus-derived systems, e.g., vectors derived from bacterial plasmids, from bacteriophage, from transposons, from yeast episomes, from insertion elements, from yeast chromosomal elements, from viruses such as baculoviruses, papova viruses, such as SV40, vaccinia viruses, adenoviruses, fowl pox viruses, pseudorabies viruses and retroviruses, and vectors derived from combinations thereof, such as those derived from plasmid and bacteriophage genetic elements, such as cosmids and phagemids. The expression systems may contain control regions that regulate as well as engender expression. Generally, any system or vector suitable to maintain, propagate or express polynucleotides to produce a polypeptide in a host may be used. The appropriate nucleotide sequence may be inserted into an expression system by any of a variety of well-known and routine techniques, such as, for example, those set forth in Sambrook et al., *MOLECULAR CLONING, A LABORATORY MANUAL* (*supra*).

For secretion of the translated protein into the lumen of the endoplasmic reticulum, into the periplasmic space or into the extracellular environment, appropriate secretion signals may be incorporated into the desired polypeptide. These signals may be endogenous to the polypeptide or they may be heterologous signals.

If the hFrizzled-6 polypeptide is to be expressed for use in screening assays, generally, it is preferred that the polypeptide be produced at the surface of the cell. In this event, the cells may be harvested prior to use in the screening assay. If hFrizzled-6 polypeptide is secreted into the medium, the medium can be recovered in order to recover and purify the polypeptide; if produced intracellularly, the cells must first be lysed before the polypeptide is recovered.

hFrizzled-6 polypeptides can be recovered and purified from recombinant cell cultures by well-known methods including ammonium sulfate or ethanol precipitation, acid extraction, anion or cation exchange chromatography, phosphocellulose chromatography, hydrophobic interaction chromatography, affinity chromatography, hydroxylapatite chromatography and lectin chromatography. Most preferably, high performance liquid chromatography is employed for purification. Well known techniques for refolding proteins may be employed to regenerate active conformation when the polypeptide is denatured during isolation and or purification.

### Diagnostic Assays

This invention also relates to the use of hFrizzled-6 polynucleotides for use as diagnostic reagents. Detection of a mutated form of hFrizzled-6 gene associated with a dysfunction will provide a diagnostic tool that can add to or define a diagnosis of a disease or susceptibility to a disease which results from under-expression, over-expression or altered expression of hFrizzled-6. Individuals carrying mutations in the hFrizzled-6 gene may be detected at the DNA level by a variety of techniques.

Nucleic acids for diagnosis may be obtained from a subject's cells, such as from blood, urine, saliva, tissue biopsy or autopsy material. The genomic DNA may be used directly for detection or may be amplified enzymatically by using PCR or other amplification techniques prior to analysis. RNA or cDNA may also be used in similar fashion. Deletions and insertions can be detected by a change in size of the amplified product in comparison to the normal genotype. Point mutations can be identified by hybridizing amplified DNA to labeled hFrizzled-6 nucleotide sequences. Perfectly matched sequences can be distinguished from mismatched duplexes by RNase digestion or by differences in melting temperatures. DNA sequence differences may also be detected by alterations in electrophoretic mobility of DNA fragments in gels, with or without denaturing agents, or by direct DNA sequencing. See, e.g., Myers et al., *Science* (1985) 230:1242. Sequence changes at specific locations may also be revealed by nuclease protection assays, such as RNase and S1 protection or the chemical cleavage method. See Cotton et al., *Proc Natl Acad Sci USA* (1985) 85: 4397-4401. In another embodiment, an array of oligonucleotides probes comprising hFrizzled-6 nucleotide sequence or fragments thereof can be constructed to conduct efficient screening of e.g., genetic mutations. Array technology methods are well known and have general applicability and can be used to address a variety of questions in molecular genetics including gene expression, genetic linkage, and genetic variability. (See for example: M.Chee et al., *Science*, Vol 274,

pp 610-613 (1996)).

The diagnostic assays offer a process for diagnosing or determining a susceptibility to infections such as bacterial, fungal, protozoan and viral infections, particularly infections caused by HIV-1 or HIV-2; pain; cancers: diabetes, obesity; anorexia; bulimia; asthma; Parkinson's disease: acute heart failure; hypotension; hypertension; urinary retention; osteoporosis; angina pectoris; myocardial infarction; ulcers; asthma; allergies; benign prostatic hypertrophy; and psychotic and neurological disorders, including anxiety, schizophrenia, manic depression, delirium, dementia, severe mental retardation and dyskinesias, such as Huntington's disease or Gilles de la Tourette's syndrome through detection of mutation in the hFrizzled-6 gene by the methods described.

In addition, infections such as bacterial, fungal, protozoan and viral infections, particularly infections caused by HIV-1 or HIV-2; pain; cancers; diabetes, obesity; anorexia; bulimia; asthma; Parkinson's disease; acute heart failure; hypotension; hypertension; urinary retention; osteoporosis; angina pectoris; myocardial infarction; ulcers; asthma; allergies; benign prostatic hypertrophy; and psychotic and neurological disorders, including anxiety, schizophrenia, manic depression, delirium, dementia, severe mental retardation and dyskinesias, such as Huntington's disease or Gilles de la Tourette's syndrome, can be diagnosed by methods comprising determining from a sample derived from a subject an abnormally decreased or increased level of hFrizzled-6 polypeptide or hFrizzled-6 mRNA. Decreased or increased expression can be measured at the RNA level using any of the methods well known in the art for the quantitation of polynucleotides, such as, for example, PCR, RT-PCR, RNase protection, Northern blotting and other hybridization methods. Assay techniques that can be used to determine levels of a protein, such as an hFrizzled-6, in a sample derived from a host are well-known to those of skill in the art. Such assay methods include radioimmunoassays, competitive-binding assays, Western Blot analysis and ELISA assays.

Thus in another aspect, the present invention relates to a diagnostic kit for a disease or susceptibility to a disease, particularly infections such as bacterial, fungal, protozoan and viral infections, particularly infections caused by HIV-1 or HIV-2; pain; cancers; diabetes, obesity; anorexia; bulimia; asthma; Parkinson's disease; acute heart failure; hypotension; hypertension; urinary retention; osteoporosis; angina pectoris; myocardial infarction; ulcers; asthma; allergies; benign prostatic hypertrophy; and psychotic and neurological disorders, including anxiety, schizophrenia, manic depression, delirium, dementia, severe mental retardation and dyskinesias, such as Huntington's disease or Gilles de la Tourette's syndrome, which comprises:

- (a) a hFrizzled-6 polynucleotide, preferably the nucleotide sequence of SEQ ID NO: 1, or a fragment thereof;
- (b) a nucleotide sequence complementary to that of (a);
- (c) a hFrizzled-6 polypeptide, preferably the polypeptide of SEQ ID NO: 2, or a fragment thereof; or
- (d) an antibody to a hFrizzled-6 polypeptide, preferably to the polypeptide of SEQ ID NO: 2.

It will be appreciated that in any such kit, (a), (b), (c) or (d) may comprise a substantial component.

### Chromosome Assays

The nucleotide sequences of the present invention are also valuable for chromosome identification. The sequence is specifically targeted to and can hybridize with a particular location on an individual human chromosome. The mapping of relevant sequences to chromosomes according to the present invention is an important first step in correlating those sequences with gene associated disease. Once a sequence has been mapped to a precise chromosomal location, the physical position of the sequence on the chromosome can be correlated with genetic map data. Such data are found, for example, in V. McKusick, Mendelian Inheritance in Man (available on line through Johns Hopkins University Welch Medical Library). The relationship between genes and diseases that have been mapped to the same chromosomal region are then identified through linkage analysis (coinheritance of physically adjacent genes). The differences in the cDNA or genomic sequence between affected and unaffected individuals can also be determined. If a mutation is observed in some or all of the affected individuals but not in any normal individuals, then the mutation is likely to be the causative agent of the disease.

### Antibodies

The polypeptides of the invention or their fragments or analogs thereof, or cells expressing them can also be used as immunogens to produce antibodies immunospecific for the hFrizzled-6 polypeptides. The term "immunospecific" means that the antibodies have substantially greater affinity for the polypeptides of the invention than their affinity for other related polypeptides in the prior art.

Antibodies generated against the hFrizzled-6 polypeptides can be obtained by administering the polypeptides or epitope-bearing fragments, analogs or cells to an animal, preferably a nonhuman, using routine protocols. For preparation of monoclonal antibodies, any technique which provides antibodies produced by continuous cell line cultures

can be used. Examples include the hybridoma technique (Kohler, G. and Milstein, C., *Nature* (1975) 256:495-497), the trioma technique, the human B-cell hybridoma technique (Kozbor *et al.*, *Immunology Today* (1983) 4:72) and the EBV-hybridoma technique (Cole *et al.*, MONOCLONAL ANTIBODIES AND CANCER THERAPY, pp. 77-96, Alan R. Liss, Inc., 1985).

Techniques for the production of single chain antibodies (U.S. Patent No. 4,946,778) can also be adapted to produce single chain antibodies to polypeptides of this invention. Also, transgenic mice, or other organisms including other mammals, may be used to express humanized antibodies.

The above-described antibodies may be employed to isolate or to identify clones expressing the polypeptide or to purify the polypeptides by affinity chromatography.

Antibodies against hFrizzled-6 polypeptides may also be employed to treat infections such as bacterial, fungal, protozoan and viral infections, particularly infections caused by HIV-1 or HIV-2; pain; cancers; diabetes, obesity; anorexia; bulimia; asthma; Parkinson's disease; acute heart failure; hypotension; hypertension; urinary retention; osteoporosis; angina pectoris; myocardial infarction; ulcers; asthma; allergies; benign prostatic hypertrophy; and psychotic and neurological disorders, including anxiety, schizophrenia, manic depression, delirium, dementia, severe mental retardation and dyskinesias, such as Huntington's disease or Gilles de la Tourette's syndrome, among others.

### Vaccines

Another aspect of the invention relates to a method for inducing an immunological response in a mammal which comprises inoculating the mammal with hFrizzled-6 polypeptide, or a fragment thereof, adequate to produce antibody and/or T cell immune response to protect said animal from infections such as bacterial, fungal, protozoan and viral infections, particularly infections caused by HIV-1 or HIV-2; pain; cancers; diabetes, obesity; anorexia; bulimia; asthma; Parkinson's disease; acute heart failure; hypotension; hypertension; urinary retention; osteoporosis; angina pectoris; myocardial infarction; ulcers; asthma; allergies; benign prostatic hypertrophy; and psychotic and neurological disorders, including anxiety, schizophrenia, manic depression, delirium, dementia, severe mental retardation and dyskinesias, such as Huntington's disease or Gilles de la Tourette's syndrome, among others. Yet another aspect of the invention relates to a method of inducing immunological response in a mammal which comprises, delivering hFrizzled-6 polypeptide via a vector directing expression of hFrizzled-6 polynucleotide *in vivo* in order to induce such an immunological response to produce antibody to protect said animal from diseases.

Further aspect of the invention relates to an immunological/vaccine formulation (composition) which, when introduced into a mammalian host, induces an immunological response in that mammal to a hFrizzled-6 polypeptide wherein the composition comprises a hFrizzled-6 polypeptide or hFrizzled-6 gene. The vaccine formulation may further comprise a suitable carrier. Since hFrizzled-6 polypeptide may be broken down in the stomach, it is preferably administered parenterally (including subcutaneous, intramuscular, intravenous, intradermal etc. injection). Formulations suitable for parenteral administration include aqueous and non-aqueous sterile injection solutions which may contain anti-oxidants, buffers, bacteriostats and solutes which render the formulation isotonic with the blood of the recipient; and aqueous and non-aqueous sterile suspensions which may include suspending agents or thickening agents. The formulations may be presented in unit-dose or multi-dose containers, for example, sealed ampoules and vials and may be stored in a freeze-dried condition requiring only the addition of the sterile liquid carrier immediately prior to use. The vaccine formulation may also include adjuvant systems for enhancing the immunogenicity of the formulation, such as oil-in-water systems and other systems known in the art. The dosage will depend on the specific activity of the vaccine and can be readily determined by routine experimentation.

### Screening Assays

The hFrizzled-6 polypeptide of the present invention may be employed in a screening process for compounds which bind the receptor and which activate (agonists) or inhibit activation of (antagonists) the receptor polypeptide of the present invention. Thus, polypeptides of the invention may also be used to assess the binding of small molecule substrates and ligands in, for example, cells, cell-free preparations, chemical libraries, and natural product mixtures. These substrates and ligands may be natural substrates and ligands or may be structural or functional mimetics. See Coligan *et al.*, *Current Protocols in Immunology* 1(2):Chapter 5 (1991).

hFrizzled-6 polypeptides are responsible for many biological functions, including many pathologies. Accordingly, it is desirable to find compounds and drugs which stimulate hFrizzled-6 on the one hand and which can inhibit the function of hFrizzled-6 on the other hand. In general, agonists are employed for therapeutic and prophylactic purposes for such conditions as infections such as bacterial, fungal, protozoan and viral infections, particularly infections caused by HIV-1 or HIV-2; pain; cancers; diabetes, obesity; anorexia; bulimia; asthma; Parkinson's disease; acute heart failure; hypotension; hypertension; urinary retention; osteoporosis; angina pectoris; myocardial infarction; ulcers; asthma; allergies; benign prostatic hypertrophy; and psychotic and neurological disorders, including anxiety, schizophrenia, manic

depression, delirium, dementia, severe mental retardation and dyskinesias, such as Huntington's disease or Gilles de la Tourette's syndrome. Antagonists may be employed for a variety of therapeutic and prophylactic purposes for such conditions as infections such as bacterial, fungal, protozoan and viral infections, particularly infections caused by HIV-1 or HIV-2; pain; cancers; diabetes, obesity; anorexia; bulimia; asthma; Parkinson's disease; acute heart failure; hypotension; hypertension; urinary retention; osteoporosis; angina pectoris; myocardial infarction; ulcers; asthma; allergies; benign prostatic hypertrophy; and psychotic and neurological disorders, including anxiety, schizophrenia, manic depression, delirium, dementia, severe mental retardation and dyskinesias, such as Huntington's disease or Gilles de la Tourette's syndrome.

In general, such screening procedures involve producing appropriate cells which express the receptor polypeptide of the present invention on the surface thereof. Such cells include cells from mammals, yeast, *Drosophila* or *E. coli*. Cells expressing the receptor (or cell membrane containing the expressed receptor) are then contacted with a test compound to observe binding, or stimulation or inhibition of a functional response.

One screening technique includes the use of cells which express the receptor of this invention (for example, transfected CHO cells) in a system which measures extracellular pH or intracellular calcium changes caused by receptor activation. In this technique, compounds may be contacted with cells expressing the receptor polypeptide of the present invention. A second messenger response, e.g., signal transduction, pH changes, or changes in calcium level, is then measured to determine whether the potential compound activates or inhibits the receptor.

Another method involves screening for receptor inhibitors by determining inhibition or stimulation of receptor-mediated cAMP and/or adenylate cyclase accumulation. Such a method involves transfecting a eukaryotic cell with the receptor of this invention to express the receptor on the cell surface. The cell is then exposed to potential antagonists in the presence of the receptor of this invention. The amount of cAMP accumulation is then measured. If the potential antagonist binds the receptor, and thus inhibits receptor binding, the levels of receptor-mediated cAMP, or adenylate cyclase, activity will be reduced or increased. Another method for detecting agonists or antagonists for the receptor of the present invention is the yeast based technology as described in U.S. Patent 5,482,835.

The assays may simply test binding of a candidate compound wherein adherence to the cells bearing the receptor is detected by means of a label directly or indirectly associated with the candidate compound or in an assay involving competition with a labeled competitor. Further, these assays may test whether the candidate compound results in a signal generated by activation of the receptor, using detection systems appropriate to the cells bearing the receptor at their surfaces. Inhibitors of activation are generally assayed in the presence of a known agonist and the effect on activation by the agonist by the presence of the candidate compound is observed.

Further, the assays may simply comprise the steps of mixing a candidate compound with a solution containing a hFrizzled-6 polypeptide to form a mixture, measuring hFrizzled-6 activity in the mixture, and comparing the hFrizzled-6 activity of the mixture to a standard.

The hFrizzled-6 cDNA, protein and antibodies to the protein may also be used to configure assays for detecting the effect of added compounds on the production of hFrizzled-6 mRNA and protein in cells. For example, an ELISA may be constructed for measuring secreted or cell associated levels of hFrizzled-6 protein using monoclonal and polyclonal antibodies by standard methods known in the art, and this can be used to discover agents which may inhibit or enhance the production of hFrizzled-6 (also called antagonist or agonist, respectively) from suitably manipulated cells or tissues. Standard methods for conducting screening assays are well understood in the art.

Examples of potential hFrizzled-6 antagonists include antibodies or, in some cases, oligonucleotides or proteins which are closely related to the ligand of the hFrizzled-6, e.g., a fragment of the ligand, or small molecules which bind to the receptor but do not elicit a response, so that the activity of the receptor is prevented.

Thus in another aspect, the present invention relates to a screening kit for identifying agonists, antagonists, ligands, receptors, substrates, enzymes, etc. for hFrizzled-6 polypeptides; or compounds which decrease or enhance the production of hFrizzled-6 polypeptides, which comprises:

- (a) a hFrizzled-6 polypeptide, preferably that of SEQ ID NO:2;
- (b) a recombinant cell expressing a hFrizzled-6 polypeptide, preferably that of SEQ ID NO:2;
- (c) a cell membrane expressing a hFrizzled-6 polypeptide; preferably that of SEQ ID NO: 2; or
- (d) antibody to a hFrizzled-6 polypeptide, preferably that of SEQ ID NO: 2.

It will be appreciated that in any such kit, (a), (b), (c) or (d) may comprise a substantial component.

### Prophylactic and Therapeutic Methods

This invention provides methods of treating an abnormal conditions related to both an excess of and insufficient amounts of hFrizzled-6 activity.

If the activity of hFrizzled-6 is in excess, several approaches are available. One approach comprises administering

to a subject an inhibitor compound (antagonist) as hereinabove described along with a pharmaceutically acceptable carrier in an amount effective to inhibit activation by blocking binding of ligands to the hFrizzled-6, or by inhibiting a second signal, and thereby alleviating the abnormal condition.

In another approach, soluble forms of hFrizzled-6 polypeptides still capable of binding the ligand in competition with endogenous hFrizzled-6 may be administered. Typical embodiments of such competitors comprise fragments of the hFrizzled-6 polypeptide.

In still another approach, expression of the gene encoding endogenous hFrizzled-6 can be inhibited using expression blocking techniques. Known such techniques involve the use of antisense sequences, either internally generated or separately administered. See, for example, O'Connor, *J Neurochem* (1991) 56:560 in Oligodeoxynucleotides as Antisense Inhibitors of Gene Expression, CRC Press, Boca Raton, FL (1988). Alternatively, oligonucleotides which form triple helices with the gene can be supplied. See, for example, Lee *et al.*, *Nucleic Acids Res* (1979) 6:3073; Cooney *et al.*, *Science* (1988) 241:456; Dervan *et al.*, *Science* (1991) 251:1360. These oligomers can be administered *per se* or the relevant oligomers can be expressed *in vivo*.

For treating abnormal conditions related to an under-expression of hFrizzled-6 and its activity, several approaches are also available. One approach comprises administering to a subject a therapeutically effective amount of a compound which activates hFrizzled-6, i.e., an agonist as described above, in combination with a pharmaceutically acceptable carrier, to thereby alleviate the abnormal condition. Alternatively, gene therapy may be employed to effect the endogenous production of hFrizzled-6 by the relevant cells in the subject. For example, a polynucleotide of the invention may be engineered for expression in a replication defective retroviral vector, as discussed above. The retroviral expression construct may then be isolated and introduced into a packaging cell transduced with a retroviral plasmid vector containing RNA encoding a polypeptide of the present invention such that the packaging cell now produces infectious viral particles containing the gene of interest. These producer cells may be administered to a subject for engineering cells *in vivo* and expression of the polypeptide *in vivo*. For overview of gene therapy, see Chapter 20, *Gene Therapy and other Molecular Genetic-based Therapeutic Approaches*, (and references cited therein) in Human Molecular Genetics, T Strachan and A P Read, BIOS Scientific Publishers Ltd (1996).

### Formulation and Administration

Peptides, such as the soluble form of hFrizzled-6 polypeptides, and agonists and antagonist peptides or small molecules, may be formulated in combination with a suitable pharmaceutical carrier. Such formulations comprise a therapeutically effective amount of the polypeptide or compound, and a pharmaceutically acceptable carrier or excipient. Such carriers include but are not limited to, saline, buffered saline, dextrose, water, glycerol, ethanol, and combinations thereof. Formulation should suit the mode of administration, and is well within the skill of the art. The invention further relates to pharmaceutical packs and kits comprising one or more containers filled with one or more of the ingredients of the aforementioned compositions of the invention.

Polypeptides and other compounds of the present invention may be employed alone or in conjunction with other compounds, such as therapeutic compounds.

Preferred forms of systemic administration of the pharmaceutical compositions include injection, typically by intravenous injection. Other injection routes, such as subcutaneous, intramuscular, or intraperitoneal, can be used. Alternative means for systemic administration include transmucosal and transdermal administration using penetrants such as bile salts or fusidic acids or other detergents. In addition, if properly formulated in enteric or encapsulated formulations, oral administration may also be possible. Administration of these compounds may also be topical and/or localized, in the form of salves, pastes, gels and the like.

The dosage range required depends on the choice of peptide, the route of administration, the nature of the formulation, the nature of the subject's condition, and the judgment of the attending practitioner. Suitable dosages, however, are in the range of 0.1-100 µg/kg of subject. Wide variations in the needed dosage, however, are to be expected in view of the variety of compounds available and the differing efficiencies of various routes of administration. For example, oral administration would be expected to require higher dosages than administration by intravenous injection. Variations in these dosage levels can be adjusted using standard empirical routines for optimization, as is well understood in the art.

Polypeptides used in treatment can also be generated endogenously in the subject, in treatment modalities often referred to as "gene therapy" as described above. Thus, for example, cells from a subject may be engineered with a polynucleotide, such as a DNA or RNA, to encode a polypeptide *ex vivo*, and for example, by the use of a retroviral plasmid vector. The cells are then introduced into the subject.

### Examples

The examples below are carried out using standard techniques, which are well known and routine to those of skill

in the art, except where otherwise described in detail. The examples illustrate, but do not limit the invention.

#### Example 1: Mammalian Cell Expression

The receptors of the present invention are expressed in either human embryonic kidney 293 (HEK293) cells or adherent dhfr CHO cells. To maximize receptor expression, typically all 5' and 3' untranslated regions (UTRs) are removed from the receptor cDNA prior to insertion into a pCDN or pCDNA3 vector. The cells are transfected with individual receptor cDNAs by lipofectin and selected in the presence of 400 mg/ml G418. After 3 weeks of selection, individual clones are picked and expanded for further analysis. HEK293 or CHO cells transfected with the vector alone serve as negative controls. To isolate cell lines stably expressing the individual receptors, about 24 clones are typically selected and analyzed by Northern blot analysis. Receptor mRNAs are generally detectable in about 50% of the G418-resistant clones analyzed.

#### Example 2 Ligand bank for binding and functional assays.

A bank of over 200 putative receptor ligands has been assembled for screening. The bank comprises: transmitters, hormones and chemokines known to act via a human seven transmembrane (7TM) receptor; naturally occurring compounds which may be putative agonists for a human 7TM receptor, non-mammalian, biologically active peptides for which a mammalian counterpart has not yet been identified; and compounds not found in nature, but which activate 7TM receptors with unknown natural ligands. This bank is used to initially screen the receptor for known ligands, using both functional (i.e. calcium, cAMP, microphysiometer, oocyte electrophysiology, etc, see below) as well as binding assays.

#### Example 3: Ligand Binding Assays

Ligand binding assays provide a direct method for ascertaining receptor pharmacology and are adaptable to a high throughput format. The purified ligand for a receptor is radiolabeled to high specific activity (50-2000 Ci/mmol) for binding studies. A determination is then made that the process of radiolabeling does not diminish the activity of the ligand towards its receptor. Assay conditions for buffers, ions, pH and other modulators such as nucleotides are optimized to establish a workable signal to noise ratio for both membrane and whole cell receptor sources. For these assays, specific receptor binding is defined as total associated radioactivity minus the radioactivity measured in the presence of an excess of unlabeled competing ligand. Where possible, more than one competing ligand is used to define residual nonspecific binding.

#### Example 4: Functional Assay in Xenopus Oocytes

Capped RNA transcripts from linearized plasmid templates encoding the receptor cDNAs of the invention are synthesized in vitro with RNA polymerases in accordance with standard procedures. In vitro transcripts are suspended in water at a final concentration of 0.2 mg/ml. Ovarian lobes are removed from adult female toads, Stage V defolliculated oocytes are obtained, and RNA transcripts (10 ng/oocyte) are injected in a 50 nl bolus using a microinjection apparatus. Two electrode voltage clamps are used to measure the currents from individual Xenopus oocytes in response to agonist exposure. Recordings are made in Ca<sup>2+</sup> free Barth's medium at room temperature. The Xenopus system can be used to screen known ligands and tissue/cell extracts for activating ligands.

#### Example 5: Microphysiometric Assays

Activation of a wide variety of secondary messenger systems results in extrusion of small amounts of acid from a cell. The acid formed is largely as a result of the increased metabolic activity required to fuel the intracellular signaling process. The pH changes in the media surrounding the cell are very small but are detectable by the CYTOSENSOR microphysiometer (Molecular Devices Ltd., Menlo Park, CA). The CYTOSENSOR is thus capable of detecting the activation of a receptor which is coupled to an energy utilizing intracellular signaling pathway such as the G-protein coupled receptor of the present invention.

#### Example 6: Extract/Cell Supernatant Screening

A large number of mammalian receptors exist for which there remains, as yet, no cognate activating ligand (agonist). Thus, active ligands for these receptors may not be included within the ligands banks as identified to date. Accordingly, the 7TM receptor of the invention is also functionally screened (using calcium, cAMP, microphysiometer,

oocyte electrophysiology, etc., functional screens) against tissue extracts to identify natural ligands. Extracts that produce positive functional responses can be sequentially subfractionated until an activating ligand is isolated identified.

#### Example 7: Calcium and cAMP Functional Assays

7TM receptors which are expressed in HEK 293 cells have been shown to be coupled functionally to activation of PLC and calcium mobilization and/or cAMP stimulation or inhibition. Basal calcium levels in the HEK 293 cells in receptor-transfected or vector control cells were observed to be in the normal, 100 nM to 200 nM, range. HEK 293 cells expressing recombinant receptors are loaded with fura 2 and in a single day > 150 selected ligands or tissue/cell extracts are evaluated for agonist induced calcium mobilization. Similarly, HEK 293 cells expressing recombinant receptors are evaluated for the stimulation or inhibition of cAMP production using standard cAMP quantitation assays. Agonists presenting a calcium transient or cAMP fluctuation are tested in vector control cells to determine if the response is unique to the transfected cells expressing receptor.

All publications, including but not limited to patents and patent applications, cited in this specification are herein incorporated by reference as if each individual publication were specifically and individually indicated to be incorporated by reference herein as though fully set forth.



Annex to the description

## SEQUENCE LISTING

5

## (1) GENERAL INFORMATION

10

(i) APPLICANT: SmithKline Beecham Corporation

(ii) TITLE OF THE INVENTION: A Human 7-TM Receptor Similar  
to Murine Frizzled-6 gene

15

(iii) NUMBER OF SEQUENCES: 4

(iv) CORRESPONDENCE ADDRESS:

20

(A) ADDRESSEE: SmithKline Beecham, Corporate Intellectual  
Property

(B) STREET: Two New Horizons Court

(C) CITY: Brentford

25

(D) STATE: Middlesex

(E) COUNTRY: United Kingdom

(F) ZIP: TW8 9EP

30

(v) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Diskette

(B) COMPUTER: IBM Compatible

(C) OPERATING SYSTEM: DOS

35

(D) SOFTWARE: FastSEQ for Windows Version 2.0

(vi) CURRENT APPLICATION DATA:

40

(A) APPLICATION NUMBER: TO BE ASSIGNED

(B) FILING DATE: 09-DEC-1997

(C) CLASSIFICATION: UNKNOWN

45

(vii) PRIOR APPLICATION DATA:

(A) APPLICATION NUMBER: 60/048,240

(B) FILING DATE: 02-JUN-1997

50

(viii) ATTORNEY/AGENT INFORMATION:

(A) NAME: CONNELL, Anthony Christopher

(B) REGISTRATION NUMBER: 5630

55

(C) REFERENCE/DOCKET NUMBER: GH70060

## (ix) TELECOMMUNICATION INFORMATION:

- (A) TELEPHONE: +44 1279 644 395  
 (B) TELEFAX: +44 181 975 6294  
 (C) TELEX:

## (2) INFORMATION FOR SEQ ID NO:1:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 2880 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: cDNA

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

```

25  CCGAGCCGTC TCAGGTCCCT GGGGGGAACG GTGGGTTAGA CGGGGACGGG AAGGGACAGC    60
    GGCCTTCGAC CGCCCCCGGA GTAATTGACC CAGGACTCAT TTTCAGGAAA GCCTGAAAAT    120
    GAGTAAATA GTGAAATGAG GAATTTGAAC ATTTTATCTT TGGATGGGGA TCTTCTGAGG    180
    ATGCAAAGAG TGATTCATCC AAGCCATGTG GTAAAATCAG GAATTTGAAG AAAATGGAGA    240
30  TGTTTACATT TTTGTTGACG TGTATTTTTC TACCCCTCCT AAGAGGGCAC AGTCTCTTCA    300
    CCTGTGAACC AATTACTGTT CCCAGATGTA TGAAAATGGC CTACAACATG ACGTTTTTCC    360
    CTAATCTGAT GGGTCATTAT GACCAGAGTA TTGCCGCGGT GGAAATGGAG CATTTTCTTC    420
    CTCTCGCAA TCTGGAATGT TCACCAAACA TTGAAACTTT CCTCTGCAA GCATTTGTAC    480
35  CAACCTGCAT AGAACAAATT CATGTGGTTC CACCTTGTCG TAAACTTTGT GAGAAAGTAT    540
    ATTCTGATTG CAAAAAATTA ATTGACACTT TTGGGATCCG ATGGCCTGAG GAGCTTGAAT    600
    GTGACAGATT ACAATACTGT GATGAGACTG TTCCTGTAAC TTTTGATCCA CACACAGAAT    660
    TTCTTGGTCC TCAGAAGGAA ACAGAACAAG TCCAAAGAGA CATTTGGATT TGGTGTCCAA    720
40  GGCATCTTAA GACTTCTGGG GGACAAGGAT ATAAGTTTCT GGAATTGAC CAGTGTGCGC    780
    CTCCATGCCC CAACATGTAT TTTAAAAGTG ATGAGCTAGA GTTTGCAAAA AGTTTTATTG    840
    GAACAGTTTC AATATTTTGT CTTTGTGCAA CTCTGTTCAC ATTCCTTACT TTTTAAATTG    900
    ATGTTAGAAG ATTCAGATAC CCAGAGAGAC CAATTATATA TTAATCTGTC TGTTACAGCA    960
45  TTGTATCTCT TATGTACTTC ATTGGATTTT TGCTGGGCGA TAGCACAGCC TGCAATAAGG   1020
    CAGATGAGAA GCTAGAACTT GGTGACACTG TTGTCTAGG CTCTCAAAAT AAGGCTTGCA   1080
    CCGTTTTTGT CATGCTTTTG TATTTTTTCA CAATGGCTGG CACTGTGTGG TGGGTGATTC   1140
50  TTACCATTAC TTGGTTCTTA GCTGCAGGAA GAAAATGGAG TTGTGAAGCC ATCGAGCAAA   1200
    AAGCAGTGTG GTTTCATGCT GTTGATGGG GAACACCAAG TTTCTGACT GTTATGCTTC   1260
    TTGCTCTGAA CAAAGTTGAA GGAGACAACA TTAGTGGAGT TCGCTTTGTT GGCCTTTATG   1320
    ACCTGGATGC TTCTCGCTAC TTTGTACTCT TGCCACTGTG CCTTTGTGTG TTTGTTGGGC   1380
55  TCTCTCTTCT TTTAGCTGGC ATTATTTTCT TAAATCATGT TCGACAAGTC ATACAACATG   1440

```

5 ATGGCCGGAA CCAAGAAAAA CTAAAGAAAT TTATGATTCTG AATTGGAGTC TTCAGCGGCT 1500  
 TGTATCTTGT GCCATTAGTG ACACTTCTCG GATGTTACGT CTATGAGCAA GTGAACAGGA 1560  
 10 TTACCTGGGA GATAACTTGG GTCTCTGATC ATTGTCGTCA GTACCATATC CCATGTCCTT 1620  
 ATCAGGCAAA AGCAAAAGCT CGACCAGAAT TGGCTTTATT TATGATAAAA TACCTGATGA 1680  
 CATTAATTGT TGGCATCTCT GCTGTCTTCT GGGTTGGAAG CAAAAAGACA TGCACAGAAT 1740  
 GGGCTGGGTT TTTTAAACGA AATCGCGGGA GAGATCCAAT CAGTGAAAGT CGAAGAGTAC 1800  
 15 TACAGGAATC ATGTGAGTTT TTCTTAAAGC ACAATTCTAA AGTTAAACAC AAAAAGAAGC 1860  
 ACTATAAACC GAGTTCACGC AAGCTGAAGG TCATTTCCAA ATCCATGGGA ACCAGCACAG 1920  
 GAGCTACAGC AAATCATGGC ACTTCTGCAG TAGCAATTAC TAGCCATGAT TACCTAGGAC 1980  
 AAGAACTTT GACAGAAATC CAAACCTCAC CAGAAACATC AATGAGAGAG GTGAAAGCGG 2040  
 20 ACGGAGCTAG CACCCCCAGG TTAAGAGAAC AGGACTGTGG TGAACCTGCC TCGCCAGCAG 2100  
 CATCCATCTC CAGACTCTCT GGGGAACAGG TCGACGGGAA GGGCCAGGCA GGCAGTGTAT 2160  
 CTGAAAGTGC GCGGAGTGAA GGAAGGATTA GTCCAAAGAG TGATATTACT GACACTGSCC 2220  
 TGGCACAGAG CAACAATTTG CAGGTCCCCA GTTCTTCAGA ACCAAGCAGC CTCAAAGGTT 2280  
 CCACATCTCT GCTTGTTTAC CCAGTTTCAG GAGTGAGAAA AGAGCAGGGA GGTGGTTGTC 2340  
 ATTCAGATAC TTGAAGAACA TTTTCTCTCG TTACTCAGAA GCAAATTTGT GTTACACTGG 2400  
 AAGTGACCTA TGCACTGTTT TGTAAGAATC ACTGTTACGT TCTTCTTTTG CACTTAAAGT 2460  
 25 TGCATTGCCT ACTGTTATAC TGGAAAAAAT AGAGTTCAAG AATAATATGA CTCATTTCAC 2520  
 ACAAAGGTTA ATGACAACAA TATACCTGAA AACAGAAATG TGCAGGTAA TAATATTTTT 2580  
 TTAATAGTGT GGGAGGACAG AGTTAGAGGA ATCTTCCTTT TCTATTTATG AAGATTCTAC 2640  
 TCTTGGAAG AGTATTTTAA GATGTACTAT GCTATTTTAC TTTTGTGATA TAAAATCAAG 2700  
 30 ATATTTCTTT GCTGAAGTAT TTAAATCTTA TCCTTGTATC TTTTATACA TATTTGAAAA 2760  
 TAAGCTTATA TGTATTTGAA CTTTTTTGAA ATCCTATTCC AGTATTTTTA TCATGCTATT 2820  
 GTGATATTTT AGCACTTTGG TAGCTTTTAC ACTGAAATTT CTAAGAAAAT TGTAATAATAG 2880

35 (2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

- 40 (A) LENGTH: 706 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

45 (ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

50 Met Glu Met Phe Thr Phe Leu Leu Thr Cys Ile Phe Leu Pro Leu Leu  
 1 5 10 15  
 Arg Gly His Ser Leu Phe Thr Cys Glu Pro Ile Thr Val Pro Arg Cys  
 20 25 30  
 55 Met Lys Met Ala Tyr Asn Met Thr Phe Phe Pro Asn Leu Met Gly His

|    |   |   |     |     |    |
|----|---|---|-----|-----|----|
|    | 35  |   | 40  |     | 45 |
|    | Tyr Asp Gln Ser Ile Ala   | Ala Val Glu Met Glu His Phe Leu Pro Leu |     |     |    |
| 5  | 50  | 55                                      | 60  |     |    |
|    | Ala Asn Leu Glu Cys Ser Pro Asn Ile Glu Thr Phe Leu Cys Lys Ala |   |     |     |    |
|    | 65  | 70                                      | 75  | 80  |    |
|    | Phe Val Pro Thr Cys Ile Glu Gln Ile His Val Val Pro Pro Cys Arg |   |     |     |    |
| 10 | 85  | 90                                      | 95  |     |    |
|    | Lys Leu Cys Glu Lys Val Tyr Ser Asp Cys Lys Lys Leu Ile Asp Thr |   |     |     |    |
|    | 100   | 105                                     | 110 |     |    |
|    | Phe Gly Ile Arg Trp Pro Glu Glu Leu Glu Cys Asp Arg Leu Gln Tyr |   |     |     |    |
| 15 | 115   | 120                                     | 125 |     |    |
|    | Cys Asp Glu Thr Val Pro Val Thr Phe Asp Pro His Thr Glu Phe Leu |   |     |     |    |
|    | 130   | 135                                     | 140 |     |    |
|    | Gly Pro Gln Lys Glu Thr Glu Gln Val Gln Arg Asp Ile Gly Phe Trp |   |     |     |    |
| 20 | 145   | 150                                     | 155 | 160 |    |
|    | Cys Pro Arg His Leu Lys Thr Ser Gly Gly Gln Gly Tyr Lys Phe Leu |   |     |     |    |
|    | 165   | 170                                     | 175 |     |    |
|    | Gly Ile Asp Gln Cys Ala Pro Pro Cys Pro Asn Met Tyr Phe Lys Ser |   |     |     |    |
| 25 | 180   | 185                                     | 190 |     |    |
|    | Asp Glu Leu Glu Phe Ala Lys Ser Phe Ile Gly Thr Val Ser Ile Phe |   |     |     |    |
|    | 195   | 200                                     | 205 |     |    |
|    | Cys Leu Cys Ala Thr Leu Phe Thr Phe Leu Thr Phe Leu Ile Asp Val |   |     |     |    |
| 30 | 210   | 215                                     | 220 |     |    |
|    | Arg Arg Phe Arg Tyr Pro Glu Arg Pro Ile Ile Tyr Tyr Ser Val Cys |   |     |     |    |
|    | 225   | 230                                     | 235 | 240 |    |
|    | Tyr Ser Ile Val Ser Leu Met Tyr Phe Ile Gly Phe Leu Leu Gly Asp |   |     |     |    |
| 35 | 245   | 250                                     | 255 |     |    |
|    | Ser Thr Ala Cys Asn Lys Ala Asp Glu Lys Leu Glu Leu Gly Asp Thr |   |     |     |    |
|    | 260   | 265                                     | 270 |     |    |
|    | Val Val Leu Gly Ser Gln Asn Lys Ala Cys Thr Val Leu Phe Met Leu |   |     |     |    |
| 40 | 275   | 280                                     | 285 |     |    |
|    | Leu Tyr Phe Phe Thr Met Ala Gly Thr Val Trp Trp Val Ile Leu Thr |   |     |     |    |
|    | 290   | 295                                     | 300 |     |    |
|    | Ile Thr Trp Phe Leu Ala Ala Gly Arg Lys Trp Ser Cys Glu Ala Ile |   |     |     |    |
| 45 | 305   | 310                                     | 315 | 320 |    |
|    | Glu Gln Lys Ala Val Trp Phe His Ala Val Ala Trp Gly Thr Pro Ser |   |     |     |    |
|    | 325   | 330                                     | 335 |     |    |
|    | Phe Leu Thr Val Met Leu Leu Ala Leu Asn Lys Val Glu Gly Asp Asn |   |     |     |    |
| 50 | 340   | 345                                     | 350 |     |    |
|    | Ile Ser Gly Val Arg Phe Val Gly Leu Tyr Asp Leu Asp Ala Ser Arg |   |     |     |    |
|    | 355   | 360                                     | 365 |     |    |
| 55 |   |   |     |     |    |
|    | Tyr Phe Val Leu Leu Pro Leu Cys Leu Cys Val Phe Val Gly Leu Ser |   |     |     |    |

|    |     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |  |  |  |
|----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|--|--|--|
|    | 370 |     | 375 |     | 380 |     |     |     |     |     |     |     |     |     |     |     |  |  |  |
|    | Leu | Leu | Leu | Ala | Gly | Ile | Ile | Ser | Leu | Asn | His | Val | Arg | Gln | Val | Ile |  |  |  |
| 5  | 385 |     |     |     |     | 390 |     |     |     |     | 395 |     |     |     | 400 |     |  |  |  |
|    | Gln | His | Asp | Gly | Arg | Asn | Gln | Glu | Lys | Leu | Lys | Lys | Phe | Met | Ile | Arg |  |  |  |
|    |     |     |     |     | 405 |     |     |     |     | 410 |     |     |     |     | 415 |     |  |  |  |
|    | Ile | Gly | Val | Phe | Ser | Gly | Leu | Tyr | Leu | Val | Pro | Leu | Val | Thr | Leu | Leu |  |  |  |
| 10 |     |     |     | 420 |     |     |     |     | 425 |     |     |     |     | 430 |     |     |  |  |  |
|    | Gly | Cys | Tyr | Val | Tyr | Glu | Gln | Val | Asn | Arg | Ile | Thr | Trp | Glu | Ile | Thr |  |  |  |
|    |     |     |     | 435 |     |     |     |     | 440 |     |     |     |     | 445 |     |     |  |  |  |
|    | Trp | Val | Ser | Asp | His | Cys | Arg | Gln | Tyr | His | Ile | Pro | Cys | Pro | Tyr | Gln |  |  |  |
| 15 |     |     |     | 450 |     |     |     |     | 455 |     |     |     |     | 460 |     |     |  |  |  |
|    | Ala | Lys | Ala | Lys | Ala | Arg | Pro | Glu | Leu | Ala | Leu | Phe | Met | Ile | Lys | Tyr |  |  |  |
|    | 465 |     |     |     |     | 470 |     |     |     |     | 475 |     |     |     | 480 |     |  |  |  |
| 20 | Leu | Met | Thr | Leu | Ile | Val | Gly | Ile | Ser | Ala | Val | Phe | Trp | Val | Gly | Ser |  |  |  |
|    |     |     |     |     | 485 |     |     |     |     |     | 490 |     |     |     | 495 |     |  |  |  |
|    | Lys | Lys | Thr | Cys | Thr | Glu | Trp | Ala | Gly | Phe | Phe | Lys | Arg | Asn | Arg | Gly |  |  |  |
|    |     |     |     | 500 |     |     |     |     | 505 |     |     |     |     | 510 |     |     |  |  |  |
| 25 | Arg | Asp | Pro | Ile | Ser | Glu | Ser | Arg | Arg | Val | Leu | Gln | Glu | Ser | Cys | Glu |  |  |  |
|    |     |     |     | 515 |     |     |     |     | 520 |     |     |     |     | 525 |     |     |  |  |  |
|    | Phe | Phe | Leu | Lys | His | Asn | Ser | Lys | Val | Lys | His | Lys | Lys | Lys | His | Tyr |  |  |  |
|    |     |     |     | 530 |     |     |     |     | 535 |     |     |     |     | 540 |     |     |  |  |  |
| 30 | Lys | Pro | Ser | Ser | Arg | Lys | Leu | Lys | Val | Ile | Ser | Lys | Ser | Met | Gly | Thr |  |  |  |
|    | 545 |     |     |     |     | 550 |     |     |     |     | 555 |     |     |     | 560 |     |  |  |  |
|    | Ser | Thr | Gly | Ala | Thr | Ala | Asn | His | Gly | Thr | Ser | Ala | Val | Ala | Ile | Thr |  |  |  |
|    |     |     |     |     | 565 |     |     |     |     |     | 570 |     |     |     | 575 |     |  |  |  |
| 35 | Ser | His | Asp | Tyr | Leu | Gly | Gln | Glu | Thr | Leu | Thr | Glu | Ile | Gln | Thr | Ser |  |  |  |
|    |     |     |     | 580 |     |     |     |     |     | 585 |     |     |     | 590 |     |     |  |  |  |
|    | Pro | Glu | Thr | Ser | Met | Arg | Glu | Val | Lys | Ala | Asp | Gly | Ala | Ser | Thr | Pro |  |  |  |
|    |     |     |     | 595 |     |     |     |     | 600 |     |     |     |     | 605 |     |     |  |  |  |
| 40 | Arg | Leu | Arg | Glu | Gln | Asp | Cys | Gly | Glu | Pro | Ala | Ser | Pro | Ala | Ala | Ser |  |  |  |
|    |     |     |     | 610 |     |     |     |     | 615 |     |     |     |     | 620 |     |     |  |  |  |
|    | Ile | Ser | Arg | Leu | Ser | Gly | Glu | Gln | Val | Asp | Gly | Lys | Gly | Gln | Ala | Gly |  |  |  |
|    | 625 |     |     |     |     | 630 |     |     |     |     | 635 |     |     |     | 640 |     |  |  |  |
| 45 | Ser | Val | Ser | Glu | Ser | Ala | Arg | Ser | Glu | Gly | Arg | Ile | Ser | Pro | Lys | Ser |  |  |  |
|    |     |     |     |     | 645 |     |     |     |     |     | 650 |     |     |     | 655 |     |  |  |  |
|    | Asp | Ile | Thr | Asp | Thr | Gly | Leu | Ala | Gln | Ser | Asn | Asn | Leu | Gln | Val | Pro |  |  |  |
|    |     |     |     | 660 |     |     |     |     |     | 665 |     |     |     | 670 |     |     |  |  |  |
| 50 | Ser | Ser | Ser | Glu | Pro | Ser | Ser | Leu | Lys | Gly | Ser | Thr | Ser | Leu | Leu | Val |  |  |  |
|    |     |     |     | 675 |     |     |     |     | 680 |     |     |     |     | 685 |     |     |  |  |  |
|    | His | Pro | Val | Ser | Gly | Val | Arg | Lys | Glu | Gln | Gly | Gly | Gly | Cys | His | Ser |  |  |  |
| 55 |     |     |     | 690 |     |     |     |     | 695 |     |     |     |     | 700 |     |     |  |  |  |
|    | Asp | Thr |     |     |     |     |     |     |     |     |     |     |     |     |     |     |  |  |  |

705

5 (2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:

10 (A) LENGTH: 315 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

15 (ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

20 TCTGAACAAA GTTGAAGGAG ACAACATTAG TGGAGTTTGC TTTGTTGGCC TTTATGACCT 60  
GGATGCTTCT CGCTACTTTG TACTCTTGCC ACTGTGCCTT TNTGTGTTTG TTGGGCTCTC 120  
TCTTCTTTTA GCTGGCATT A TTTCCTTAA TCATGTTTGA CAAGTCATAC AACATGATGG 180  
25 CCGGAACCAA GAAAACTAA AGAAATTTAT GATTCTGAATT GGAGTCTTCA GCGGCTTTGT 240  
ATCTTGTGCC ATTAGTGACA CTTCTCGGAT GTTACGTCTA TGAGCAAGTG AACAGGATTA 300  
CCTGGGAGAT AACTT 315

30 (2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:

35 (A) LENGTH: 26 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

40 (ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

45 Leu Asn Lys Val Glu Gly Asp Asn Ile Ser Gly Val Cys Phe Val Gly  
1 5 10 15  
Leu Tyr Asp Leu Asp Ala Ser Arg Tyr Phe  
50 20 25

## Claims

- 55 1. An isolated polynucleotide comprising a nucleotide sequence that has at least 80% identity over its entire length to a nucleotide sequence encoding the hFrizzled-6 polypeptide of SEQ ID NO:2; or a nucleotide sequence complementary to said isolated polynucleotide.

2. The polynucleotide of claim 1 wherein said polynucleotide comprises the nucleotide sequence contained in SEQ ID NO: 1 encoding the hFrizzled-6 polypeptide of SEQ ID NO:2.

3. The polynucleotide of claim 1 wherein said polynucleotide comprises a nucleotide sequence that is at least 80% identical to that of SEQ ID NO: 1 over its entire length.

4. The polynucleotide of claim 3 which is polynucleotide of SEQ ID NO: 1.

5. The polynucleotide of claim 1 which is DNA or RNA.

6. A DNA or RNA molecule comprising an expression system, wherein said expression system is capable of producing a hFrizzled-6 polypeptide comprising an amino acid sequence, which has at least 80% identity with the polypeptide of SEQ ID NO:2 when said expression system is present in a compatible host cell.

7. A host cell comprising the expression system of claim 6.

8. A process for producing a hFrizzled-6 polypeptide comprising culturing a host of claim 7 under conditions sufficient for the production of said polypeptide and recovering the polypeptide from the culture.

9. A process for producing a cell which produces a hFrizzled-6 polypeptide thereof comprising transforming or transfecting a host cell with the expression system of claim 6 such that the host cell, under appropriate culture conditions, produces a hFrizzled-6 polypeptide.

10. A hFrizzled-6 polypeptide comprising an amino acid sequence which is at least 80% identical to the amino acid sequence of SEQ ID NO:2 over its entire length.

11. The polypeptide of claim 10 which comprises the amino acid sequence of SEQ ID NO:2.

12. An antibody immunospecific for the hFrizzled-6 polypeptide of claim 10.

13. A method for the treatment of a subject in need of enhanced activity or expression of hFrizzled-6 polypeptide of claim 10 comprising:

- (a) administering to the subject a therapeutically effective amount of an agonist to said receptor; and/or
- (b) providing to the subject an isolated polynucleotide comprising a nucleotide sequence that has at least 80% identity over its entire length to a nucleotide sequence encoding the hFrizzled-6 polypeptide of SEQ ID NO: 2; or a nucleotide sequence complementary to said isolated polynucleotide in a form so as to effect production of said receptor activity *in vivo*.

14. A method for the treatment of a subject having need to inhibit activity or expression of hFrizzled-6 polypeptide of claim 10 comprising:

- (a) administering to the subject a therapeutically effective amount of an antagonist to said receptor; and/or
- (b) administering to the subject a nucleic acid molecule that inhibits the expression of the nucleotide sequence encoding said receptor; and/or
- (c) administering to the subject a therapeutically effective amount of a polypeptide that competes with said receptor for its ligand.

15. A process for diagnosing a disease or a susceptibility to a disease in a subject related to expression or activity of hFrizzled-6 polypeptide of claim 10 in a subject comprising:

- (a) determining the presence or absence of a mutation in the nucleotide sequence encoding said hFrizzled-6 polypeptide in the genome of said subject; and/or
- (b) analyzing for the presence or amount of the hFrizzled-6 polypeptide expression in a sample derived from said subject.

16. A method for identifying agonists to hFrizzled-6 polypeptide of claim 10 comprising:

- (a) contacting a cell which produces a hFrizzled-6 polypeptide with a candidate compound; and
- (b) determining whether the candidate compound effects a signal generated by activation of the hFrizzled-6 polypeptide.

5 17. An agonist identified by the method of claim 16.

18. The method for identifying antagonists to hFrizzled-6 polypeptide of claim 10 comprising:

- 10 (a) contacting a cell which produces a hFrizzled-6 polypeptide with an agonist; and
- (b) determining whether the signal generated by said agonist is diminished in the presence of a candidate compound.

19. An antagonist identified by the method of claim 18.

15 20. A recombinant host cell produced by a method of Claim 9 or a membrane thereof expressing a hFrizzled-6 polypeptide.

20

25

30

35

40

45

50

55



(19)



Europäisches Patentamt  
European Patent Office  
Office européen des brevets



(11)

**EP 0 882 793 A3**

(12)

**EUROPEAN PATENT APPLICATION**

(88) Date of publication A3:  
17.05.2000 Bulletin 2000/20

(43) Date of publication A2:  
09.12.1998 Bulletin 1998/50

(21) Application number: 98304208.6

(22) Date of filing: 28.05.1998

(51) Int Cl.7: **C12N 15/12**, C12N 15/85,  
C12N 15/11, C12N 5/10,  
C12Q 1/02, C12Q 1/68,  
C07K 14/705, C07K 16/28,  
A61K 38/17, A61K 48/00,  
G01N 33/50

(84) Designated Contracting States:  
**AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU**  
**MC NL PT SE**  
Designated Extension States:  
**AL LT LV MK RO SI**

(30) Priority: 02.06.1997 US 48240 P  
09.12.1997 US 987289

(71) Applicant: **SMITHKLINE BEECHAM**  
**CORPORATION**  
Philadelphia Pennsylvania 19103 (US)

(72) Inventors:  
• Hu, Erding  
King of Prussia, Pennsylvania 19406 (US)  
• Zhu, Yuan  
King of Prussia, Pennsylvania 19406 (US)

(74) Representative:  
Crump, Julian Richard John et al  
FJ Cleveland,  
40-43 Chancery Lane  
London WC2A 1JQ (GB)

(54) **A human 7-tm receptor similar to murine frizzled-6 gene**

(57) hFrizzled-6 polypeptides and polynucleotides and methods for producing such polypeptides by recombinant techniques are disclosed. Also disclosed are methods for utilizing hFrizzled-6 polypeptides and polynucleotides in the design of protocols for the treatment of infections such as bacterial, fungal, protozoan and viral infections, particularly infections caused by HIV-1 or HIV-2; pain; cancers; diabetes; obesity; anorexia; bulimia; asthma; Parkinson's disease; acute heart fail-

ure; hypotension; hypertension; urinary retention; osteoporosis; angina pectoris; myocardial infarction; ulcers; asthma; allergies; benign prostatic hypertrophy; and psychotic and neurological disorders, including anxiety, schizophrenia, manic depression, delirium, dementia, severe mental retardation and dyskinesias, such as Huntington's disease or Gilles de la Tourette's syndrome, among others and diagnostic assays for such conditions.

**EP 0 882 793 A3**



European Patent  
Office

# PARTIAL EUROPEAN SEARCH REPORT

Application Number

which under Rule 45 of the European Patent Convention EP 98 30 4208  
shall be considered, for the purposes of subsequent  
proceedings, as the European search report

| DOCUMENTS CONSIDERED TO BE RELEVANT  |  |                                  |   |
|--|--|----------------------------------|---|
| Category   | Citation of document with indication, where appropriate, of relevant passages  | Relevant to claim                | CLASSIFICATION OF THE APPLICATION (Int.Cl.6)  |
| X,D  | WANG Y ET AL: "A LARGE FAMILY OF PUTATIVE TRANSMEMBRANE RECEPTORS HOMOLOGOUS TO THE PRODUCT OF THE DROSOPHILA TISSUE POLARITY GENE FRIZZLED"<br>JOURNAL OF BIOLOGICAL CHEMISTRY,<br>vol. 271, no. 8,<br>23 February 1996 (1996-02-23), pages<br>4468-4476, XP002054778<br>* the whole document *<br>& "AC U43319"<br>EMBL DATABASE,<br>13 January 1996 (1996-01-13),<br>Heidelberg<br>* the whole document * | 1,3,5-7,<br>10,12                | C12N15/12<br>C12N15/85<br>C12N15/11<br>C12N5/10<br>C12Q1/02<br>C12Q1/68<br>C07K14/705<br>C07K16/28<br>A61K38/17<br>A61K48/00<br>G01N33/50 |
| X  | HOANG B ET AL: "PRIMARY STRUCTURE AND TISSUE DISTRIBUTION OF FRZB, A NOVEL PROTEIN RELATED TO DROSOPHILA FRIZZLED, SUGGEST A ROLE IN SKELETAL MORPHOGENESIS"<br>JOURNAL OF BIOLOGICAL CHEMISTRY,<br>vol. 271, no. 42,<br>18 October 1996 (1996-10-18), pages<br>26131-26137, XP002054777<br>* the whole document *   | 17,19                            | TECHNICAL FIELDS<br>SEARCHED (Int.Cl.6)<br><br>C07K<br>C12N<br>C12Q<br>A61K<br>G01N   |
| -/--   |  |                                  |   |
| <b>INCOMPLETE SEARCH</b>   |  |                                  |   |
| <p>The Search Division considers that the present application, or one or more of its claims, does/do not comply with the EPC to such an extent that a meaningful search into the state of the art cannot be carried out, or can only be carried out partially, for these claims.</p> <p>Claims searched completely :</p> <p>Claims searched incompletely :</p> <p>Claims not searched :</p> <p>Reason for the limitation of the search:</p> <p>Although claims 13,14 are directed to a method of treatment of the human/animal body (Article 52(4) EPC), the search has been carried out and based on the alleged effects of the compound/composition.</p> |  |                                  |   |
| Place of search  |  | Date of completion of the search | Examiner  |
| THE HAGUE  |  | 16 September 1998                | Kania, T  |
| <p>CATEGORY OF CITED DOCUMENTS</p> <p>X : particularly relevant if taken alone<br/>Y : particularly relevant if combined with another document of the same category<br/>A : technological background<br/>O : non-written disclosure<br/>P : intermediate document</p> <p>T : theory or principle underlying the invention<br/>E : earlier patent document, but published on, or after the filing date<br/>D : document cited in the application<br/>L : document cited for other reasons<br/>&amp; : member of the same patent family, corresponding document</p>  |  |                                  |   |

EPO FORM 1503 03.82 (P04C07)



European Patent  
Office

## PARTIAL EUROPEAN SEARCH REPORT

Application Number  
EP 98 30 4208

| DOCUMENTS CONSIDERED TO BE RELEVANT |   |                   | CLASSIFICATION OF THE APPLICATION (Int.Cl.6) |
|-------------------------------------|---|-------------------|--|
| Category                            | Citation of document with indication, where appropriate, of relevant passages   | Relevant to claim |  |
| A                                   | HILLIER L. ET AL.: "The WashU-Merck EST Project, AC AA064897"<br>EMBL DATABASE,<br>26 September 1996 (1996-09-26),<br>XP002074749<br>Heidelberg<br>* the whole document *   | 1-7,<br>10-12     |  |
| A                                   | ADAMS M. ET AL.: "AC AA385032"<br>EMBL DATABASE, 18 April 1997 (1997-04-18),<br>XP002074750<br>Heidelberg<br>* the whole document *   | 1-7,<br>10-12     |  |
| A                                   | WANG Y K ET AL: "A novel human homologue of the Drosophila frizzled wnt receptor gene binds wingless protein and is in the Williams syndrome deletion at 7q11.23."<br>HUMAN MOLECULAR GENETICS, (1997 MAR) 6 (3) 465-72. JOURNAL CODE: BRC. ISSN: 0964-6906., XP002074751<br>* the whole document * | 1-20              | TECHNICAL FIELDS SEARCHED (Int.Cl.6)         |
| A                                   | HE X ET AL: "A member of the Frizzled protein family mediating axis induction by Wnt-5A."<br>SCIENCE, (1997 MAR 14) 275 (5306) 1652-4. JOURNAL CODE: UJ7. ISSN: 0036-8075., XP002074752<br>* the whole document *   | 1-20              |  |
| P,X                                 | TOKUHARA M ET AL: "Molecular cloning of human Frizzled-6."<br>BIOCHEMICAL AND BIOPHYSICAL RESEARCH COMMUNICATIONS, (1998 FEB 13) 243 (2) 622-7. JOURNAL CODE: 9Y8. ISSN: 0006-291X., XP002074753<br>* the whole document *  | 1-20              |  |
|                                     | ---   | ---               |  |
|                                     | -/--  |                   |  |

EPO FORM 1503 03.82 (P04C10)

European Patent  
Office

## PARTIAL EUROPEAN SEARCH REPORT

Application Number  
EP 98 30 4208

| DOCUMENTS CONSIDERED TO BE RELEVANT |  |                   | CLASSIFICATION OF THE APPLICATION (Int.Cl.6) |
|-------------------------------------|--|-------------------|--|
| Category                            | Citation of document with indication, where appropriate, of relevant passages  | Relevant to claim |  |
| T                                   | WO 97 39357 A (UNIV LELAND STANFORD JUNIOR<br>;UNIV JOHNS HOPKINS (US))<br>23 October 1997 (1997-10-23)<br>* the whole document *<br>----- | 1-20              |  |
|                                     |  |                   | TECHNICAL FIELDS<br>SEARCHED (Int.Cl.6)      |
|                                     |  |                   |  |
|                                     |  |                   |  |

EPO FORM 1503 03/82 (P04C10)

**ANNEX TO THE EUROPEAN SEARCH REPORT  
ON EUROPEAN PATENT APPLICATION NO.**

EP 98 30 4208

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report.  
The members are as contained in the European Patent Office EDP file on  
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

16-09-1998

| Patent document<br>cited in search report | Publication<br>date | Patent family<br>member(s) | Publication<br>date |
|---|---------------------|----------------------------|---------------------|
| WO 9739357 A                              | 23-10-1997          | AU 2664797 A               | 07-11-1997          |
| -----                                     |                     |                            |                     |
|   |                     |                            |                     |
|   |                     |                            |                     |
|   |                     |                            |                     |
|   |                     |                            |                     |
|   |                     |                            |                     |
|   |                     |                            |                     |
|   |                     |                            |                     |
|   |                     |                            |                     |
|   |                     |                            |                     |
|   |                     |                            |                     |

EPO FORM P0458

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82

